arXiv:1202.3347v1 [math-ph] 15 Feb 2012

Indecomposable representations and oscillator realizations of the exceptional Lie algebra G_2

Hua-Jun Huang, You-Ning Li, and Dong Ruan [*]

Department of Physics, Tsinghua University, Beijing 100084, P.R. China

February 16, 2012

Abstract

In this paper various representations of the exceptional Lie algebra G_2 are investigated in a purely algebraic manner, and multi-boson/multi-fermion realizations are obtained. Matrix elements of the master representation, which is defined on the space of the universal enveloping algebra of G_2, are explicitly determined. From this master representation, different indecomposable representations defined on invariant subspaces or quotient spaces with respect to these invariant subspaces are discussed. Especially, the elementary representations of G_2 are investigated in detail, and the corresponding six-boson realization is given. After obtaining explicit forms of all twelve extremal vectors of the elementary representation with the highest weight $\mathbf{\Lambda}$, all representations with their respective highest weights related to $\mathbf{\Lambda}$ are systematically discussed. For one of these representations the corresponding five-boson realization is constructed. Moreover, a new three-fermion realization from the fundamental representation (0,1) of G_2 is constructed also.

---

[*] dongruan@tsinghua.edu.cn

I INTRODUCTION

The exceptional Lie algebra G_2 has many physical applications. The first successful application is in atomic spectroscopy, where Racah[1] used G_2 to classify the states of atomic f^n configurations according to the Lie algebra chain $so(7) \supset G_2 \supset so(3)$ and to simplify the calculations of the matrix elements of the Coulomb interaction. Soon afterwards, Racah's idea and method was directly applied to nuclear physics[2] to classify the orbital states of nuclear f^n configurations in Russell-Saunders coupling according to the Lie algebra chain $u(7) \supset so(7) \supset G_2 \supset so(3)$.[3,4] Morrison *et al.*[5] discussed the interacting boson model including s, d, g, i bosons by use of $u(28) \supset u(7) \supset so(7) \supset G_2 \supset so(3)$. In particle physics, G_2 symmetry is drawing more attention recently and is the topic of even more works.[6,7,8,9,10] These applications motivate us to study G_2.

Considerable research has been done investigating G_2. Irreducible matrix representations of G_2 have been obtained by embedding G_2 into A_6,[11] and in the $G_2 \supset su(3)$ basis[12] as well as the $G_2 \supset so(4)$ basis,[13] respectively. The Casimir operators of degrees 2 and 6 of G_2 with its generators written in terms of the subalgebra A_2 were obtained.[14] In recent years, some works on G_2 have been put forward such as new types of representations with solitary and edge-minimal bases,[15] twist quantization,[16] and exceptional quantum subgroups.[17] However, much less is known about indecomposable representations of G_2. The indecomposable representations (i.e. reducible but not completely reducible representations) have been found useful in physics for a long time.[18,19,20,21,22,23,24] Many authors have investigated the indecomposable representations of various algebras: Lorentz algebra,[25,26] Euclidean algebra,[27,28] Poincaré algebra,[29] Lie superalgebras,[22,30] quantum groups,[31,32] the Diamond Lie algebra,[33] and so on. Especially, Gruber and co-workers[34,35,36,37] have studied in detail the indecomposable representations of A_1, A_2, and D_2 on the spaces of their respective universal enveloping algebras and on the quotient spaces with respect to the invariant subspaces. Recently the indecomposable representations appeared to be topical in the representation theory of the nonlinear Lie algebras.[38,39] The indecomposable representations can also be used to construct various boson realizations of the (nonlinear) Lie algebras.[40,41,42]

In this paper we will first investigate the master representation of G_2 on the space of its universal enveloping algebra, and various representations which are defined on different invariant subspaces and on quotient spaces with respect to different ideals. Then from these representations we will construct different oscillator realizations in terms of six bosons, five bosons, and three fermions, respectively.

This paper is arranged as follows: In Sec. II, the general procedure of obtaining inhomogeneous boson realizations (IHBRs)[39] from the master representation or from the induced representations is briefly reviewed, which is applicable to any Lie algebra. In Sec. III, explicit forms of the matrix elements of the master representation ρ of G_2 are determined, and its various indecomposable representations and the corresponding IHBRs are discussed respectively. In Sec. IV, the matrix elements of the elementary representation $d_{\mathbf{\Lambda}}$ with $\mathbf{\Lambda}$ being the highest weight, as an induced representation of ρ, are calculated. Then a six-boson realization from $d_{\mathbf{\Lambda}}$ is constructed.

In Sec. V, all twelve extremal vectors of $d_{\mathbf{\Lambda}}$ are determined. They may define various invariant subspaces which carry various elementary sub-representations. Then we investigate two series of representations on different quotient spaces. Different cases of each series are discussed, a five-boson realization is obtained from one of them. Moveover, to find realizations with fewer boson or fermion operators, we manage to construct a new three-fermion realization from the fundamental representation (0,1) as an induced representation of $d_{\mathbf{\Lambda}}$. A summary and some discussions are given in the final section.

In this paper, $\mathbb{N}$ denotes the set of positive integers, $\mathbb{Z}^+$ the set of non-negative integers, and $\mathbb{C}$ the set of complex numbers.

II THE GENERAL PROCEDURE

Consider a Lie algebra $\mathfrak{g}$ with generators T_i $(i = 1, 2, \ldots, m, m = \dim \mathfrak{g})$. According to the Poincare-Birkhoff-Witt theorem, a basis for the universal enveloping algebra $\mathfrak{U}(\mathfrak{g})$ of $\mathfrak{g}$ can be naturally chosen as the following set of ordered elements:

$$\{X(i_1, i_2, \ldots, i_m) = T_1^{i_1} T_2^{i_2} \ldots T_m^{i_m} | i_1, i_2, \ldots, i_m \in \mathbb{Z}^+\}. \tag{1}$$

We call $\{X(i_1, i_2, \ldots, i_m)\}$ the PBW basis of $\mathfrak{g}$. The left action of the generators T_a upon this PBW basis (1) gives

$$\rho(T_a) X(i_1, i_2, \ldots, i_m) = T_a T_1^{i_1} T_2^{i_2} \ldots T_m^{i_m} = \sum_{i'_1, \ldots, i'_m} \rho(T_a)^{i'_1, i'_2, \ldots, i'_m}_{i_1, i_2, \ldots, i_m} X(i'_1, i'_2, \ldots, i'_m). \tag{2}$$

Here we obtain a representation ρ of $\mathfrak{g}$ on $\mathfrak{U}(\mathfrak{g})$, called the master representation. The matrix elements $\rho(T_a)^{i'_1 i'_2 \ldots i'_m}_{i_1 i_2 \ldots i_m}$ can be explicitly determined by the commutation relations of $\mathfrak{g}$. Let I be a left ideal of $\mathfrak{U}(\mathfrak{g})$. On the quotient space $\mathfrak{U}(\mathfrak{g})/I$, ρ may induce a representation, and different choices of I enable us to obtain various representations of $\mathfrak{g}$.

Now we can construct the IHBR of $\mathfrak{g}$ from the master representation ρ on $\mathfrak{U}(\mathfrak{g})$ or from the representations on $\mathfrak{U}(\mathfrak{g})/I$. Since the matrix elements $\rho(T_a)^{i'_1, i'_2, \ldots, i'_m}_{i_1, i_2, \ldots, i_m}$ in Eq. (2) are related to m independent parameters $i_1, i_2, \ldots, i_m$, we need m pairs of independent boson operators $\{a_i^\dagger, a_i\}$ $(i = 1, 2, \ldots, m)$ to define a boson Fock space $\mathcal{F}_b$ with basis

$$\mathcal{F}_b : \{|i_1, i_2, \ldots, i_m\rangle \equiv {a_1^\dagger}^{i_1} {a_2^\dagger}^{i_2} \ldots {a_m^\dagger}^{i_m} |0\rangle \, | i_1, i_2, \ldots, i_m \in \mathbb{Z}^+\}, \tag{3}$$

where $|0\rangle$ is the vacuum state of $\mathcal{F}_b$, and $a_i|0\rangle = 0$. These boson operators satisfy the usual boson commutation relations:

$$[a_i, a_j^\dagger] = \delta_{ij}, \quad [a_i, a_j] = [a_i^\dagger, a_j^\dagger] = 0, \tag{4}$$

$$\hat{n}_i \equiv a_i^\dagger a_i, \quad [\hat{n}_i, a_j] = -\delta_{ij} a_j, \quad [\hat{n}_i, a_j^\dagger] = \delta_{ij} a_j^\dagger, \tag{5}$$

with $\hat{n}_i$ being the particle number operator of the ith boson.

Then the mapping $\Phi : \mathfrak{U}(\mathfrak{g}) \longrightarrow \mathcal{F}_b$ or $\Phi : \mathfrak{U}(\mathfrak{g})/I \longrightarrow \mathcal{F}_b$ defined by

$$\Phi(X(i_1, i_2, \ldots, i_s)) = |i_1, i_2, \ldots, i_s\rangle, \quad s \leq m \tag{6}$$

is an automorphism to $\mathfrak{U}(\mathfrak{g})$ or $\mathfrak{U}(\mathfrak{g})/I$. Let

$$\Gamma(T_i) = \Phi\rho(T_i)\Phi^{-1}. \tag{7}$$

It is obvious that $\Gamma(T_i)$ satisfy the same commutation relations as the generators T_i of $\mathfrak{g}$, i.e.,

$$[\Gamma(T_i), \Gamma(T_j)] = \Gamma([T_i, T_j]). \tag{8}$$

Therefore, Eq. (7) defines a Fock representation of $\mathfrak{g}$ on $\mathcal{F}_b$:

$$\Gamma(T_a)|i_1, i_2, \ldots, i_m\rangle = \sum_{i'_1, \ldots, i'_m} \rho(T_a)^{i'_1, i'_2, \ldots, i'_m}_{i_1, i_2, \ldots, i_m} |i'_1, i'_2, \ldots, i'_m\rangle. \tag{9}$$

Note that the matrix elements $\rho(T_a)^{i'_1, i'_2, \ldots, i'_m}_{i_1, i_2, \ldots, i_m}$ in Eq. (9) are just the same as those in Eq. (2). It is easy to obtain the corresponding IHBR of $\mathfrak{g}$ with the help of the following formulas:

$$\begin{aligned} a^\dagger_k|\ldots, i_k, \ldots\rangle &= |\ldots, i_k + 1, \ldots\rangle, \\ a_k|\ldots, i_k, \ldots\rangle &= i_k|\ldots, i_k - 1, \ldots\rangle, \\ \hat{n}_k|\ldots, i_k, \ldots\rangle &= i_k|\ldots, i_k, \ldots\rangle. \end{aligned} \tag{10}$$

III THE MASTER REPRESENTATION OF G_2

G_2 is a fourteen-dimensional simple Lie algebra of rank two.[43] Its root system Σ has twelve roots $\{\pm\alpha_i | i = 1, 2, \ldots, 6\}$. Since the Cartan subalgebra of G_2 is two-dimensional, the six positive roots in Σ may be chosen as:

$$\begin{aligned} &\alpha_1 = (\frac{1}{4}, -\frac{\sqrt{3}}{4}), \quad \alpha_2 = (\frac{1}{4}, -\frac{1}{4\sqrt{3}}), \quad \alpha_3 = (\frac{1}{2}, 0), \\ &\alpha_4 = (\frac{1}{4}, \frac{1}{4\sqrt{3}}), \quad \alpha_5 = (\frac{1}{4}, \frac{\sqrt{3}}{4}), \quad \alpha_6 = (0, \frac{1}{2\sqrt{3}}), \end{aligned} \tag{11}$$

where α_1, α_6 are two simple roots.

The Cartan-Weyl basis of G_2 reads

$$\{E_{\pm\alpha_1}, E_{\pm\alpha_2}, E_{\pm\alpha_3}, E_{\pm\alpha_4}, E_{\pm\alpha_5}, E_{\pm\alpha_6}, H_1, H_2\}. \tag{12}$$

In this paper, we will use the convenient notations $E_{\pm i} \equiv E_{\pm\alpha_i}$, which are shown in Fig. 1. The subset $\{H_1, H_2\}$ forms the Cartan subalgebra $\mathfrak{h}$ of G_2.

The commutation relations of G_2 are

$$[H_i, E_\alpha] \;=\; \alpha^{(i)} E_\alpha, \quad \text{with } \alpha^{(i)} \equiv \alpha(H_i),\ i = 1, 2; \tag{13}$$

$$[E_\alpha, E_\beta] \;=\; \begin{cases} \sum_i \alpha^{(i)} H_i, & \text{if } \alpha+\beta=0, \\ N_{\alpha,\beta} E_{\alpha+\beta}, & \text{if } \alpha+\beta \in \Sigma, \end{cases} \tag{14}$$

where coefficients $N_{\alpha,\beta}$ are respectively

$$N_{6,1} = N_{6,4} = N_{4,2} = N_{1,5} = \frac{1}{2\sqrt{2}}, \quad N_{6,2} = \frac{1}{\sqrt{6}} \tag{15}$$

with the relations $N_{\alpha,\beta} = -N_{-\alpha,-\beta}$.

The PBW basis of the universal enveloping algebra Ω of G_2 can be chosen as the following set of ordered elements

$$\Omega : \{X(m_1, \ldots, k_2) \equiv E_{-1}^{m_1} E_{-2}^{m_2} E_{-3}^{m_3} E_{-4}^{m_4} E_{-5}^{m_5} E_{-6}^{m_6} E_1^{n_1} E_2^{n_2} E_3^{n_3} E_4^{n_4} E_5^{n_5} E_6^{n_6} H_1^{k_1} H_2^{k_2} \\ | m_i, n_i, k_1, k_2 \in \mathbb{Z}^+, i = 1, 2, \ldots, 6\}. \tag{16}$$

In this paper we call the order from E_{-1} to H_2 in Eq. (16)“the standard order”, and let **1** denote the identity operator, which is with all parameters simulataneously being zero, i.e., $\mathbf{1} = X(0, 0, \ldots, 0)$. Before further computation, it is useful to derive the following general commutation relations between any two Cartan-Weyl elements of a Lie algebra $\mathfrak{g}$:

$$[H_i, E_\alpha^n] = n\alpha^{(i)} E_\alpha^n, \tag{17}$$

$$[E_\alpha, E_{-\alpha}^n] = \sum_{i=1}^{2} n\alpha^{(i)} E_{-\alpha}^{n-1} H_i + \frac{1}{2} n(n-1)(\alpha, -\alpha) E_{-\alpha}^{n-1}, \tag{18}$$

$$[E_\alpha, E_\beta^n] = \sum_m C_n^m \left(\prod_{i=1}^{m} N_{\alpha+(i-1)\beta,\beta} \right) E_\beta^{n-m} E_{\alpha+m\beta}, \tag{19}$$

where $n, m \in \mathbb{N}$, $m \le n$ and $m \le 3$, and $C_n^m \equiv \frac{n!}{m!(n-m)!}$ is the binomial coefficient.

Using above equations, the master representation ρ of G_2 can be directly calculated. The explicit matrix elements of ρ are very complicated and are put in Appendix, where we have used $X_{(mnk)}$ as abbreviation of $X(m_1, \ldots, k_2)$, X_{m_i+1} as abbreviation of $X(\ldots, m_i + 1, \ldots)$. Notice that the matrix elements for the two



Figure 1: Cartan-Weyl basis of G_2

generators $\{H_1, H_2\}$ of $\mathfrak{h}$ are

$$\begin{cases} \rho(H_1)X_{(mnk)} = X_{k_1+1} + \left(\sum_{i=1}^{6} \alpha_i^{(1)}(n_i - m_i)\right) X_{(mnk)}, \\ \rho(H_2)X_{(mnk)} = X_{k_2+1} + \left(\sum_{i=1}^{6} \alpha_i^{(2)}(n_i - m_i)\right) X_{(mnk)}. \end{cases} \tag{20}$$

Hence any element of the PBW basis $\{X_{(mnk)}\}$ of G_2 is not the common eigenvector of $\{H_1, H_2\}$.

By making use of Eq. (10), we can obtain directly from ρ an inhomogeneous fourteen-boson realization of G_2. However, we are interested in realizations with fewer boson pairs. Thus, different representations on quotient spaces with respect to different left ideals need investigating.

In ρ the parameters k_1, k_2, n_5, n_6, $n_{\widehat{45}} \equiv n_4 + n_5$, and $n_{\widehat{15}} \equiv n_1 + n_2 + 2n_3 + n_4 + n_5$ can be increased only. This means that ρ is indecomposable in any of these parameters. Therefore, each of the subspaces V_{K_1}, V_{K_2}, V_{N_5}, V_{N_6}, $V_{N_{\widehat{45}}}$, and $V_{N_{\widehat{15}}}$ of Ω, with their respective bases being

$$\begin{array}{ll} V_{K_1}: & \{E_{-1}^{m_1} \ldots E_1^{n_1} \ldots E_5^{n_5} E_6^{n_6} H_1^{K_1+k_1} H_2^{k_2} | K_1 \in \mathbb{N}\}, \\ V_{K_2}: & \{E_{-1}^{m_1} \ldots E_1^{n_1} \ldots E_5^{n_5} E_6^{n_6} H_1^{k_1} H_2^{K_2+k_2} | K_2 \in \mathbb{N}\}, \\ V_{N_5}: & \{E_{-1}^{m_1} \ldots E_1^{n_1} \ldots E_5^{N_5+n_5} E_6^{n_6} H_1^{k_1} H_2^{k_2} | N_5 \in \mathbb{N}\}, \\ V_{N_6}: & \{E_{-1}^{m_1} \ldots E_1^{n_1} \ldots E_5^{n_5} E_6^{N_6+n_6} H_1^{k_1} H_2^{k_2} | N_6 \in \mathbb{N}\}, \\ V_{N_{\widehat{45}}}: & \{E_{-1}^{m_1} \ldots E_1^{n_1} \ldots E_5^{N_{\widehat{45}}-n_4+n_5} E_6^{n_6} H_1^{k_1} H_2^{k_2} | N_{\widehat{45}} \in \mathbb{N}\}, \\ V_{N_{\widehat{15}}}: & \{E_{-1}^{m_1} \ldots E_1^{n_1} \ldots E_5^{(N_{\widehat{15}}-n_1-n_2-2n_3-n_4)+n_5} E_6^{n_6} H_1^{k_1} H_2^{k_2} | N_{\widehat{15}} \in \mathbb{N}\}, \end{array}$$

forms an invariant subspace of Ω under the action of ρ. These subspaces carry subrepresentations induced by ρ. Furthermore, ρ can also induce the representations on the quotient spaces of Ω with respect to these invariant subspaces, and their matrix elements can be obtained by formally setting

$$\begin{array}{ll} E_{-1}^{m_1} \ldots E_1^{n_1} \ldots E_5^{n_5} E_6^{n_6} H_1^{K_1} H_2^{k_2} = 0, & \text{for } \Omega/V_{K_1}, \\ E_{-1}^{m_1} \ldots E_1^{n_1} \ldots E_5^{n_5} E_6^{n_6} H_1^{k_1} H_2^{K_2} = 0, & \text{for } \Omega/V_{K_2}, \\ E_{-1}^{m_1} \ldots E_1^{n_1} \ldots E_5^{N_5} E_6^{n_6} H_1^{k_1} H_2^{k_2} = 0, & \text{for } \Omega/V_{N_5}, \\ E_{-1}^{m_1} \ldots E_1^{n_1} \ldots E_5^{n_5} E_6^{N_6} H_1^{k_1} H_2^{k_2} = 0, & \text{for } \Omega/V_{N_6}, \\ E_{-1}^{m_1} \ldots E_1^{n_1} \ldots E_5^{N_{\widehat{45}}-n_4} E_6^{n_6} H_1^{k_1} H_2^{k_2} = 0, & \text{for } \Omega/V_{N_{\widehat{45}}}, \\ E_{-1}^{m_1} \ldots E_1^{n_1} \ldots E_5^{N_{\widehat{15}}-n_1-n_2-2n_3-n_4} E_6^{n_6} H_1^{k_1} H_2^{k_2} = 0, & \text{for } \Omega/V_{N_{\widehat{15}}} \end{array}$$

in ρ, respectively. Different values of K_1, K_2, N_5, N_6, $N_{\widehat{45}}$, and $N_{\widehat{15}}$ correspond to different representations on their respective quotient spaces. For example, if we choose $K_1 = 1$, then H_1 does not appear in the PBW basis of the quotient space $\Omega/V_{K_1=1}$, that is, its basis involves only thirteen generators of G_2. The corresponding IHBR needs thirteen independent boson pairs. Moreover, since a sum of any combination of these invariant subspaces is still an invariant subspace V of Ω, a new representation on the quotient space Ω/V can be induced. Different choices of the sum spaces determine different representations. This makes it possible to find various indecomposable representations with their respective PBW bases containing only 13, 12, 11, 10, 9, 8, 7, or 6 generators of G_2 as follows:

$$\text{13 bosons}: \quad \Omega/V_{\overline{K_1}},\ \Omega/V_{\overline{K_2}},\ \Omega/V_{\overline{N_5}},\ \Omega/V_{\overline{N_6}};$$

$$
\begin{array}{rl}
12 \text{ bosons}: & \Omega/\cup_{\overline{K}_1\overline{K}_2},\ \Omega/\cup_{\overline{K}_1\overline{N}_5},\ \Omega/\cup_{\overline{K}_1\overline{N}_6},\ \Omega/\cup_{\overline{K}_2\overline{N}_5}, \\
& \Omega/\cup_{\overline{K}_2\overline{N}_6},\ \Omega/\cup_{\overline{N}_5\overline{N}_6},\ \Omega/V_{\overline{N}_{\widehat{45}}}; \\
11 \text{ bosons}: & \Omega/\cup_{\overline{K}_1\overline{K}_2\overline{N}_5},\ \Omega/\cup_{\overline{K}_1\overline{K}_2\overline{N}_6},\ \Omega/\cup_{\overline{K}_1\overline{N}_5\overline{N}_6},\ \Omega/\cup_{\overline{K}_2\overline{N}_5\overline{N}_6}, \\
& \Omega/\cup_{\overline{K}_1\overline{N}_{\widehat{45}}},\ \Omega/\cup_{\overline{K}_2\overline{N}_{\widehat{45}}},\ \Omega/\cup_{\overline{N}_{\widehat{45}}\overline{N}_6}; \\
10 \text{ bosons}: & \Omega/\cup_{\overline{K}_1\overline{K}_2\overline{N}_5\overline{N}_6},\ \Omega/\cup_{\overline{K}_1\overline{K}_2\overline{N}_{\widehat{45}}},\ \Omega/\cup_{\overline{K}_1\overline{N}_{\widehat{45}}\overline{N}_6},\ \Omega/\cup_{\overline{K}_2\overline{N}_{\widehat{45}}\overline{N}_6}; \\
9 \text{ bosons}: & \Omega/\cup_{\overline{K}_1\overline{K}_2\overline{N}_{\widehat{45}}\overline{N}_6},\ \Omega/V_{\overline{N}_{\widehat{15}}}; \\
8 \text{ bosons}: & \Omega/\cup_{\overline{K}_1\overline{N}_{\widehat{15}}},\ \Omega/\cup_{\overline{K}_2\overline{N}_{\widehat{15}}},\ \Omega/\cup_{\overline{N}_{\widehat{15}}\overline{N}_6}; \\
7 \text{ bosons}: & \Omega/\cup_{\overline{K}_1\overline{K}_2\overline{N}_{\widehat{15}}},\ \Omega/\cup_{\overline{K}_1\overline{N}_{\widehat{15}}\overline{N}_6},\ \Omega/\cup_{\overline{K}_2\overline{N}_{\widehat{15}}\overline{N}_6}; \\
6 \text{ bosons}: & \Omega/\cup_{\overline{K}_1\overline{K}_2\overline{N}_{\widehat{15}}\overline{N}_6}\ .
\end{array}
\tag{21}
$$

Here for convenience we have used the short notations: $V_{\overline{K}_1} \equiv V_{K_1=1}$ and $\cup_{\overline{K}_1\overline{K}_2\ldots} \equiv V_{K_1=1} + V_{K_2=1} + \ldots$.

Just like ρ in Eq. (20), because of the existence of X_{k_i+1}, the basis elements in most of these representations are not the common eigenvectors of $\{H_1, H_2\}$, except for the representations on $\Omega/\cup_{\overline{K}_1\overline{K}_2}$, $\Omega/\cup_{\overline{K}_1\overline{K}_2\overline{N}_5}$, $\Omega/\cup_{\overline{K}_1\overline{K}_2\overline{N}_6}$, $\Omega/\cup_{\overline{K}_1\overline{K}_2\overline{N}_5\overline{N}_6}$, $\Omega/\cup_{\overline{K}_1\overline{K}_2\overline{N}_{\widehat{45}}}$, $\Omega/\cup_{\overline{K}_1\overline{K}_2\overline{N}_{\widehat{45}}\overline{N}_6}$, $\Omega/\cup_{\overline{K}_1\overline{K}_2\overline{N}_{\widehat{15}}}$, and $\Omega/\cup_{\overline{K}_1\overline{K}_2\overline{N}_{\widehat{15}}\overline{N}_6}$.

In the case of $\Omega/\cup_{\overline{K}_1\overline{K}_2}$ with the basis

$$\{\, \bar{X}_{(mn)} \equiv E_{-1}^{m_1} E_{-2}^{m_2} E_{-3}^{m_3} E_{-4}^{m_4} E_{-5}^{m_5} E_{-6}^{m_6} E_1^{n_1} E_2^{n_2} E_3^{n_3} E_4^{n_4} E_5^{n_5} E_6^{n_6} \,\},$$

Eq. (20) now becomes

$$
\left\{
\begin{array}{l}
\rho(H_1)\bar{X}_{(mn)} = \left(\sum_{i=1}^{6} \alpha_i^{(1)}(n_i - m_i)\right) \bar{X}_{(mn)}, \\
\rho(H_2)\bar{X}_{(mn)} = \left(\sum_{i=1}^{6} \alpha_i^{(2)}(n_i - m_i)\right) \bar{X}_{(mn)}.
\end{array}
\right.
\tag{22}
$$

This set of equations shows that the basis element $\bar{X}_{(mn)}$ in $\Omega/\cup_{\overline{K}_1\overline{K}_2}$ is the common eigenvector of $\{H_1, H_2\}$, with their respective eigenvalues being

$$
\left\{
\begin{array}{ll}
\bar{M}_1 & = \sum_{i=1}^{6} \alpha_i^{(1)}(n_i - m_i) \\
& = \frac{1}{4}(n_1 + n_2 + 2n_3 + n_4 + n_5 - m_1 - m_2 - 2m_3 - m_4 - m_5), \\
\bar{M}_2 & = \sum_{i=1}^{6} \alpha_i^{(2)}(n_i - m_i) \\
& = \frac{1}{4\sqrt{3}}(-3n_1 - n_2 + n_4 + 3n_5 + 2n_6 + 3m_1 + m_2 - m_4 - 3m_5 - 2m_6).
\end{array}
\right.
\tag{23}
$$

The vector $\bar{\mathbf{M}} \equiv (\bar{M}_1, \bar{M}_2)$ is the weight of $\bar{X}_{(mn)}$. It is obvious that the weight of $\mathbf{1}$ is $\mathbf{0}$. The weight diagram of all weights $\bar{\mathbf{M}}$ of $\Omega/\cup_{\overline{K}_1\overline{K}_2}$ is shown in Fig. 2(a).

The similar arguments about $\Omega/\cup_{\overline{K}_1\overline{K}_2}$ can also be applied to the remaining seven quotient spaces: $\Omega/\cup_{\overline{K}_1\overline{K}_2\overline{N}_5}$, $\Omega/\cup_{\overline{K}_1\overline{K}_2\overline{N}_6}$, $\Omega/\cup_{\overline{K}_1\overline{K}_2\overline{N}_5\overline{N}_6}$, $\Omega/\cup_{\overline{K}_1\overline{K}_2\overline{N}_{\widehat{45}}}$, $\Omega/\cup_{\overline{K}_1\overline{K}_2\overline{N}_{\widehat{45}}\overline{N}_6}$, $\Omega/\cup_{\overline{K}_1\overline{K}_2\overline{N}_{\widehat{15}}}$, and $\Omega/\cup_{\overline{K}_1\overline{K}_2\overline{N}_{\widehat{15}}\overline{N}_6}$. For these cases, it is obvious that not all twelve parameters m_i, n_i ($i = 1, 2, \ldots, 6$) appear. The weight diagrams of $\Omega/\cup_{\overline{K}_1\overline{K}_2\overline{N}_5}$, $\Omega/\cup_{\overline{K}_1\overline{K}_2\overline{N}_6}$, $\Omega/\cup_{\overline{K}_1\overline{K}_2\overline{N}_5\overline{N}_6}$, $\Omega/\cup_{\overline{K}_1\overline{K}_2\overline{N}_{\widehat{45}}}$, and $\Omega/\cup_{\overline{K}_1\overline{K}_2\overline{N}_{\widehat{45}}\overline{N}_6}$ are the same as that of $\Omega/\cup_{\overline{K}_1\overline{K}_2}$. However, the weight diagrams of $\Omega/\cup_{\overline{K}_1\overline{K}_2\overline{N}_{\widehat{15}}}$ and $\Omega/\cup_{\overline{K}_1\overline{K}_2\overline{N}_{\widehat{15}}\overline{N}_6}$ are different from that of $\Omega/\cup_{\overline{K}_1\overline{K}_2}$, which are shown in Fig. 2 (b) and (c) respectively.

The weight diagram of $\Omega/\cup_{\overline{K}_1\overline{K}_2\overline{N}_{\widehat{15}}}$ occupies half of the full weight lattice of G_2, i.e., its weight has an upper bound in one direction. In the case of $\Omega/\cup_{\overline{K}_1\overline{K}_2\overline{N}_{\widehat{15}}\overline{N}_6}$, there exists a highest weight vector **1** with its weight being **0**, owing to the fact that all six generators E_{-i} ($i = 1$, 2,..., 6) corresponding respectively to the six negative roots appear in the basis elements of

$$\Omega/\cup_{\overline{K}_1\overline{K}_2\overline{N}_{\widehat{15}}\overline{N}_6} : \left\{E_{-1}^{m_1}E_{-2}^{m_2}E_{-3}^{m_3}E_{-4}^{m_4}E_{-5}^{m_5}E_{-6}^{m_6}\right\}.$$

Actually, the representation on $\Omega/\cup_{\overline{K}_1\overline{K}_2\overline{N}_{\widehat{15}}\overline{N}_6}$ is a special case (with the highest weight being zero) of the elementary representations, which will be discussed in the next section.

IV THE ELEMENTARY REPRESENTATIONS OF G_2

The PBW basis (16) of G_2 can be factored into a product of three subspaces: $\Omega = \Omega_-\Omega_+\mathcal{H}$, with

$$\begin{array}{rl} \Omega_- : & \{X^-(m_1, m_2, \ldots, m_6) = E_{-1}^{m_1}E_{-2}^{m_2}\ldots E_{-6}^{m_6} | m_i \in \mathbb{Z}^+, i = 1, 2, \ldots, 6\}, \\ \Omega_+ : & \{X^+(n_1, n_2, \ldots, n_6) = E_1^{n_1}E_2^{n_2}\ldots E_6^{n_6} | n_i \in \mathbb{Z}^+, i = 1, 2, \ldots, 6\}, \\ \mathcal{H} : & \{X^h(k_1, k_2) = H_1^{k_1}H_2^{k_2} | k_1, k_2 \in \mathbb{Z}^+\}. \end{array} \qquad (24)$$



Figure 2: A dot denotes a weight of the weight diagram. (a) The weight diagram of the representations on $\Omega/\cup_{\overline{K}_1\overline{K}_2}$, $\Omega/\cup_{\overline{K}_1\overline{K}_2\overline{N}_5}$, $\Omega/\cup_{\overline{K}_1\overline{K}_2\overline{N}_6}$, $\Omega/\cup_{\overline{K}_1\overline{K}_2\overline{N}_5\overline{N}_6}$, $\Omega/\cup_{\overline{K}_1\overline{K}_2\overline{N}_{\widehat{45}}}$, and $\Omega/\cup_{\overline{K}_1\overline{K}_2\overline{N}_{\widehat{45}}\overline{N}_6}$. (b) The weight diagram of the representation on $\Omega/\cup_{\overline{K}_1\overline{K}_2\overline{N}_{\widehat{15}}}$. (c) The weight diagram of the elementary representation on $\Omega/\cup_{\overline{K}_1\overline{K}_2\overline{N}_{\widehat{15}}\overline{N}_6}$, with **1** being its highest weight vector.

If on Ω one imposes the conditions

$$\left\{\begin{array}{l} \rho(H_j)\mathbf{1} = \Lambda_j\mathbf{1}, \quad \Lambda_j \in \mathbb{C},\ j = 1, 2, \\ \rho(E_i)\mathbf{1} = 0, \quad i = 1, 2, \ldots, 6, \end{array}\right. \tag{25}$$

then one may obtain a quotient space $Z(\mathbf{\Lambda}) = \Omega/I(\mathbf{\Lambda})$, where $I(\mathbf{\Lambda})$ is the left ideal of Ω generated by $\{E_1, E_2, \ldots, E_6, H_1 - \Lambda_1\mathbf{1}, H_2 - \Lambda_2\mathbf{1}\}$. Eq. (25) shows that $\mathbf{\Lambda} \equiv (\Lambda_1, \Lambda_2)$ is the weight of the vector $\mathbf{1}$, and $\mathbf{1}$ is the highest weight vector of $Z(\mathbf{\Lambda})$. The representation on $Z(\mathbf{\Lambda})$ is called the elementary representation,[35] denoted by $d_{\mathbf{\Lambda}}$. When the highest weight $\mathbf{\Lambda} = \mathbf{0}$, the representation $d_{\mathbf{\Lambda}=\mathbf{0}}$ becomes the case of $\Omega/\cup_{\overline{K}_1\overline{K}_2\overline{N}_{\widehat{15}}\overline{N}_6}$ in the last section.

Under the conditions (25), the matrix elements of the master representation ρ (see Appendix) reduce into those of the elementary representation $d_{\mathbf{\Lambda}}$. Here, using the abbreviations: $X^-_{(m)} \equiv X^-(m_1, m_2, \ldots, m_6)$ and $X^-_{m_i+k} \equiv X^-(m_1, \ldots, m_i + k, \ldots, m_6)$ in Ω_-, the explicit matrix elements of $d_{\mathbf{\Lambda}}$ are as follows:

$$\begin{aligned}
\rho(E_{-1})X^-_{(m)} &= X^-_{m_1+1}, \\
\rho(E_{-2})X^-_{(m)} &= X^-_{m_2+1}, \\
\rho(E_{-3})X^-_{(m)} &= X^-_{m_3+1}, \\
\rho(E_{-4})X^-_{(m)} &= X^-_{m_4+1} - \frac{1}{2\sqrt{2}}m_2X^-_{m_2-1,m_3+1}, \\
\rho(E_{-5})X^-_{(m)} &= X^-_{m_5+1} + \frac{1}{2\sqrt{2}}m_1X^-_{m_1-1,m_3+1}, \\
\rho(E_{-6})X^-_{(m)} &= X^-_{m_6+1} - \frac{1}{2\sqrt{2}}m_1X^-_{m_1-1,m_2+1} - \frac{1}{\sqrt{6}}m_2X^-_{m_2-1,m_4+1} \\
&\quad + \frac{1}{8\sqrt{3}}m_2(m_2 - 1)X^-_{m_2-2,m_3+1} - \frac{1}{2\sqrt{2}}m_4X^-_{m_4-1,m_5+1}, \\
\rho(E_1)X^-_{(m)} &= \frac{1}{4}m_1\left[\Lambda_1 - \sqrt{3}\Lambda_2 - \frac{1}{2}(m_1 - 1 + m_2 + m_3 - m_5 - m_6)\right]X^-_{m_1-1} \\
&\quad + \frac{1}{2\sqrt{2}}m_2X^-_{m_2-1,m_6+1} - \frac{1}{8\sqrt{3}}m_2(m_2 - 1)X^-_{m_2-2,m_4+1} \\
&\quad + \frac{1}{48\sqrt{6}}m_2(m_2 - 1)(m_2 - 2)X^-_{m_2-3,m_3+1} \\
&\quad - \frac{1}{8}m_2m_4X^-_{m_2-1,m_4-1,m_5+1} - \frac{1}{2\sqrt{2}}m_3X^-_{m_3-1,m_5+1}, \\
\rho(E_2)X^-_{(m)} &= -\frac{1}{4\sqrt{3}}m_1m_4X^-_{m_1-1,m_2+1,m_4-1} \\
&\quad + \frac{1}{16\sqrt{6}}m_1m_4(m_4 - 1)X^-_{m_1-1,m_3+1,m_4-2} - \frac{1}{8}m_1m_5X^-_{m_1-1,m_4+1,m_5-1} \\
&\quad + \frac{1}{4\sqrt{6}}m_1m_6\left(\Lambda_2 - \frac{m_6 - 1}{4\sqrt{3}}\right)X^-_{m_1-1,m_6-1} \\
&\quad + \frac{1}{4}m_2\left[\Lambda_1 - \frac{1}{\sqrt{3}}\Lambda_2 - \frac{1}{4\sqrt{3}}(3m_1 + m_2 - 1 + 3m_3 + m_4 - m_6)\right]X^-_{m_2-1}
\end{aligned}$$

$$
\begin{aligned}
&+\frac{1}{2\sqrt{2}}m_3X^-_{m_3-1,m_4+1}+\frac{1}{\sqrt{6}}m_4X^-_{m_4-1,m_6+1}\\
&-\frac{1}{8\sqrt{3}}m_4(m_4-1)X^-_{m_4-2,m_5+1},\\
\rho(E_3)X^-_{(m)} ={}& \frac{1}{16\sqrt{6}}m_1m_4(m_4-1)X^-_{m_1-1,m_2+1,m_4-2}\\
&-\frac{1}{96\sqrt{3}}m_1m_4(m_4-1)(m_4-2)X^-_{m_1-1,m_3+1,m_4-3}\\
&-\frac{1}{16\sqrt{3}}m_1m_4m_6\left(\Lambda_2-\frac{m_6-1}{4\sqrt{3}}\right)X^-_{m_1-1,m_4-1,m_6-1}\\
&-\frac{1}{8\sqrt{2}}m_1m_5\left[\Lambda_1+\sqrt{3}\Lambda_2-\frac{1}{2}\left(m_4+m_5-1+m_6\right)\right]X^-_{m_1-1,m_5-1}\\
&-\frac{1}{48\sqrt{6}}m_2(m_2-1)(m_2-2)X^-_{m_1+1,m_2-3}\\
&+\frac{1}{8\sqrt{2}}m_2m_4\left[\Lambda_1+\frac{1}{\sqrt{3}}\Lambda_2-\frac{1}{6}\left(2m_2+m_4-3+3m_5+m_6\right)\right]X^-_{m_2-1,m_4-1}\\
&+\frac{1}{8}m_2m_5X^-_{m_2-1,m_5-1,m_6+1}\\
&+\frac{1}{192}m_2(m_2-1)m_4(m_4-1)X^-_{m_2-2,m_3+1,m_4-2}\\
&-\frac{1}{16\sqrt{6}}m_2(m_2-1)m_5X^-_{m_2-2,m_4+1,m_5-1}\\
&+\frac{1}{48}m_2(m_2-1)m_6\left(\Lambda_2\ -\frac{m_6-1}{4\sqrt{3}}\right)X^-_{m_2-2,m_6-1}\\
&+\frac{1}{2}m_3\left[\Lambda_1-\frac{1}{4}\left(m_1+m_2+m_3-1+m_4+m_5\right)\right]X^-_{m_3-1}\\
&-\frac{1}{8\sqrt{3}}m_4(m_4-1)X^-_{m_4-2,m_6+1}\\
&+\frac{1}{48\sqrt{6}}m_4(m_4-1)(m_4-2)X^-_{m_4-3,m_5+1},\\
\rho(E_4)X^-_{(m)} ={}& -\frac{1}{8\sqrt{3}}m_2(m_2-1)X^-_{m_1+1,m_2-2}\\
&+\frac{1}{24\sqrt{2}}m_2m_4(m_4-1)X^-_{m_2-1,m_3+1,m_4-2}\\
&-\frac{1}{4\sqrt{3}}m_2m_5X^-_{m_2-1,m_4+1,m_5-1}\\
&+\frac{1}{6\sqrt{2}}m_2m_6\left(\Lambda_2-\frac{m_6-1}{4\sqrt{2}}\right)X^-_{m_2-1,m_6-1}\\
&-\frac{1}{2\sqrt{2}}m_3X^-_{m_2+1,m_3-1}+\frac{1}{2\sqrt{2}}m_5X^-_{m_5-1,m_6+1}\\
&+\frac{1}{4}m_4\left[\Lambda_1+\frac{1}{\sqrt{3}}\Lambda_2-\frac{1}{6}\left(4m_2+m_4-1+3m_5+m_6\right)\right]X^-_{m_4-1},
\end{aligned}
$$

$$
\begin{aligned}
\rho(E_5)X^-_{(m)} &= \frac{1}{2\sqrt{2}}m_3X^-_{m_1+1,m_3-1} - \frac{1}{8\sqrt{3}}m_4(m_4-1)X^-_{m_2+1,m_4-2} \\
&\quad + \frac{1}{24\sqrt{6}}m_4(m_4-1)(m_4-2)X^-_{m_3+1,m_4-3} \\
&\quad + \frac{1}{4\sqrt{6}}m_4m_6\left(\Lambda_2 - \frac{m_6-1}{4\sqrt{3}}\right)X^-_{m_4-1,m_6-1} \\
&\quad + \frac{1}{4}m_5\left[\Lambda_1 + \sqrt{3}\Lambda_2 - \frac{1}{2}(m_4+m_5-1+m_6)\right]X^-_{m_5-1}, \\
\rho(E_6)X^-_{(m)} &= -\frac{1}{2\sqrt{2}}m_2X^-_{m_1+1,m_2-1} - \frac{1}{\sqrt{6}}m_4X^-_{m_2+1,m_4-1} - \frac{1}{2\sqrt{2}}m_5X^-_{m_4+1,m_5-1} \\
&\quad + \frac{1}{8\sqrt{3}}m_4(m_4-1)X^-_{m_3+1,m_4-2} + \frac{1}{2\sqrt{3}}m_6\left(\Lambda_2 - \frac{m_6-1}{4\sqrt{3}}\right)X^-_{m_6-1}, \\
\rho(H_1)X^-_{(m)} &= \left[\Lambda_1 - \frac{1}{4}(m_1+m_2+2m_3+m_4+m_5)\right]X^-_{(m)} \equiv M_1X^-_{(m)}, \\
\rho(H_2)X^-_{(m)} &= \left[\Lambda_2 + \frac{1}{4\sqrt{3}}(3m_1+m_2-m_4-3m_5-2m_6)\right]X^-_{(m)} \equiv M_2X^-_{(m)}.
\end{aligned}
\tag{26}
$$

It is clear from the last two equations of Eq. (26) that $X^-_{(m)}$ is the common eigenvector of H_1 and H_2, with their respective eigenvalues being M_1 and M_2, i.e., the weight of $X^-_{(m)}$ is $\mathbf{M} \equiv (M_1, M_2)$. The weight diagram of $d_{\mathbf{\Lambda}}$ is the same as Fig. 2(c) with the highest weight being $\mathbf{\Lambda}$ instead of $\mathbf{0}$.

By making use of Eq. (10), we can immediately obtain from the matrix elements (26) of $d_{\mathbf{\Lambda}}$ the inhomogeneous six-boson realization of G_2:

$$
\begin{aligned}
&B(E_{-1}) = a_1^\dagger, \\
&B(E_{-2}) = a_2^\dagger, \\
&B(E_{-3}) = a_3^\dagger, \\
&B(E_{-4}) = a_4^\dagger - \frac{1}{2\sqrt{2}}a_3^\dagger a_2, \\
&B(E_{-5}) = a_5^\dagger + \frac{1}{2\sqrt{2}}a_3^\dagger a_1, \\
&B(E_{-6}) = a_6^\dagger - \frac{1}{2\sqrt{2}}a_2^\dagger a_1 - \frac{1}{\sqrt{6}}a_4^\dagger a_2 + \frac{1}{8\sqrt{3}}a_3^\dagger a_2^2 - \frac{1}{2\sqrt{2}}a_5^\dagger a_4, \\
&B(E_1) = \frac{1}{4}\left[\Lambda_1 - \sqrt{3}\Lambda_2 - \frac{1}{2}(\hat{n}_1+\hat{n}_2+\hat{n}_3-\hat{n}_5-\hat{n}_6)\right]a_1 \\
&\qquad + \frac{1}{2\sqrt{2}}a_6^\dagger a_2 - \frac{1}{8\sqrt{3}}a_4^\dagger a_2^2 + \frac{1}{48\sqrt{6}}a_3^\dagger a_2^3 - \frac{1}{8}a_5^\dagger a_2a_4 - \frac{1}{2\sqrt{2}}a_5^\dagger a_3, \\
&B(E_2) = -\frac{1}{4\sqrt{3}}a_2^\dagger a_1a_4 + \frac{1}{16\sqrt{6}}a_3^\dagger a_1a_4^2 - \frac{1}{8}a_4^\dagger a_1a_5 + \frac{1}{4\sqrt{6}}\left(\Lambda_2 - \frac{1}{4\sqrt{3}}\hat{n}_6\right)a_1a_6 \\
&\qquad + \frac{1}{4}\left[\Lambda_1 - \frac{1}{\sqrt{3}}\Lambda_2 - \frac{1}{4\sqrt{3}}(3\hat{n}_1+\hat{n}_2+3\hat{n}_3+\hat{n}_4-\hat{n}_6)\right]a_2 \\
&\qquad + \frac{1}{2\sqrt{2}}a_4^\dagger a_3 + \frac{1}{\sqrt{6}}a_6^\dagger a_4 - \frac{1}{8\sqrt{3}}a_5^\dagger a_4^2,
\end{aligned}
$$

$$
\begin{aligned}
B(E_3) = {} & \frac{1}{16\sqrt{6}} a_2^\dagger a_1 a_4^2 - \frac{1}{96\sqrt{3}} a_3^\dagger a_1 a_4^3 - \frac{1}{16\sqrt{3}} \left(\Lambda_2 - \frac{1}{4\sqrt{3}} \hat{n}_6 \right) a_1 a_4 a_6 \\
& - \frac{1}{8\sqrt{2}} \left[\Lambda_1 + \sqrt{3} \Lambda_2 - \frac{1}{2} (\hat{n}_4 + \hat{n}_5 + \hat{n}_6) \right] a_1 a_5 \\
& - \frac{1}{48\sqrt{6}} a_1^\dagger a_2^3 + \frac{1}{8\sqrt{2}} \left[\Lambda_1 + \frac{1}{\sqrt{3}} \Lambda_2 - \frac{1}{6} (2\hat{n}_2 + \hat{n}_4 + 3\hat{n}_5 + \hat{n}_6) \right] a_2 a_4 \\
& + \frac{1}{8} a_6^\dagger a_2 a_5 + \frac{1}{192} a_3^\dagger a_2^2 a_4^2 - \frac{1}{16\sqrt{6}} a_4^\dagger a_2^2 a_5 + \frac{1}{48} \left(\Lambda_2 - \frac{\hat{n}_6}{4\sqrt{3}} \right) a_2^2 a_6 \\
& + \frac{1}{2} \left[\Lambda_1 - \frac{1}{4} (\hat{n}_1 + \hat{n}_2 + \hat{n}_3 + \hat{n}_4 + \hat{n}_5) \right] a_3 - \frac{1}{8\sqrt{3}} a_6^\dagger a_4^2 + \frac{1}{48\sqrt{6}} a_5^\dagger a_4^3, \\
B(E_4) = {} & - \frac{1}{8\sqrt{3}} a_1^\dagger a_2^2 + \frac{1}{24\sqrt{2}} a_3^\dagger a_2 a_4^2 - \frac{1}{4\sqrt{3}} a_4^\dagger a_2 a_5 \\
& + \frac{1}{6\sqrt{2}} \left(\Lambda_2 - \frac{1}{4\sqrt{3}} \hat{n}_6 \right) a_2 a_6 - \frac{1}{2\sqrt{2}} a_2^\dagger a_3 + \frac{1}{2\sqrt{2}} a_6^\dagger a_5 \\
& + \frac{1}{4} \left[\Lambda_1 + \frac{1}{\sqrt{3}} \Lambda_2 - \frac{1}{6} (4\hat{n}_2 + \hat{n}_4 + 3\hat{n}_5 + \hat{n}_6) \right] a_4, \\
B(E_5) = {} & \frac{1}{2\sqrt{2}} a_1^\dagger a_3 - \frac{1}{8\sqrt{3}} a_2^\dagger a_4^2 + \frac{1}{24\sqrt{6}} a_3^\dagger a_4^3 \\
& + \frac{1}{4\sqrt{6}} \left(\Lambda_2 - \frac{1}{4\sqrt{3}} \hat{n}_6 \right) a_4 a_6 + \left[\frac{1}{4} \Lambda_1 + \frac{\sqrt{3}}{4} \Lambda_2 - \frac{1}{8} (\hat{n}_4 + \hat{n}_5 + \hat{n}_6) \right] a_5, \\
B(E_6) = {} & - \frac{1}{2\sqrt{2}} a_1^\dagger a_2 - \frac{1}{\sqrt{6}} a_2^\dagger a_4 - \frac{1}{2\sqrt{2}} a_4^\dagger a_5 + \frac{1}{8\sqrt{3}} a_3^\dagger a_4^2 + \frac{1}{2\sqrt{3}} \left(\Lambda_2 - \frac{1}{4\sqrt{3}} \hat{n}_6 \right) a_6, \\
B(H_1) = {} & \Lambda_1 - \frac{1}{4} (\hat{n}_1 + \hat{n}_2 + 2\hat{n}_3 + \hat{n}_4 + \hat{n}_5), \\
B(H_2) = {} & \Lambda_2 + \frac{1}{4\sqrt{3}} (3\hat{n}_1 + \hat{n}_2 - \hat{n}_4 - 3\hat{n}_5 - 2\hat{n}_6).
\end{aligned}
\tag{27}
$$

The boson realization of H_1 and H_2 is dependent on the particle number operators $\hat{n}_i$ $(i = 1, 2, \ldots, 6)$ only. Notice that $\hat{n}_6$ and $\hat{n}_3$ disappear in $B(H_1)$ and $B(H_2)$, respectively.

V TWO SERIES OF REPRESENTATIONS ON QUOTIENT SPACES OF Ω_- INDUCED FROM THE ELEMENTARY REPRESENTATION

In this section the extremal vectors defining invariant subspaces will be calculated, and all corresponding elementary representations will be determined. On quotient spaces with respect to these invariant subspaces, two series of representations induced from $d_{\mathbf{\Lambda}}$ will be discussed in detail, which are infinite-dimensional indecomposable, infinite-dimensional irreducible or finite-dimensional irreducible. The calculation of new IHBRs will be carried out in the same manner as in the last section.

V.1 Extremal vectors and the series of elementary representations

According to the definition given in Ref. [44], a vector $Y \in \Omega_-$ is called an extremal vector of the elementary representation $d_{\mathbf{\Lambda}}$, if it satisfies

$$\left\{ \begin{array}{l} \rho(H_j)Y = M_j Y, \quad M_j \in \mathbb{C},\ j = 1,2, \\ \rho(E_i)Y = 0, \quad i = 1,2,\ldots,6. \end{array} \right. \tag{28}$$

Therefore an extremal vector Y (with its weight being $\mathbf{M}$) is the highest weight vector of the subspace

$$I_Y \equiv \Omega_- Y : \{E_{-1}^{m_1} E_{-2}^{m_2} \ldots E_{-6}^{m_6} Y | m_i \in \mathbb{Z}^+, i = 1,2,\ldots,6\},$$

which is obviously an invariant subspace (ideal) of Ω_-. The representation on I_Y, which may be induced from $d_{\mathbf{\Lambda}}$, is also an elementary representation of G_2, here denoted by $d_{\mathbf{M}}$. It is clear from Eq. (28) that the identity vector $\mathbf{1}$ is an extremal vector of $d_{\mathbf{\Lambda}}$.

Bernshtein, Gel'fand, and Gel'fand[45] gave the necessary and sufficient condition that the module $d_{\mathbf{\Lambda}}$ contains $d_{\mathbf{M}}$ (known as BGG theorem): there exist a sequence of positive roots $\gamma_1, \gamma_2, \ldots, \gamma_k$ ($k \in \mathbb{N}$), which satisfy

$$\left\{ \begin{array}{l} \mathbf{M} + \mathbf{R} = S_{\gamma_k} \ldots S_{\gamma_2} S_{\gamma_1} (\mathbf{\Lambda} + \mathbf{R}), \\ 2(S_{\gamma_{i-1}} \ldots S_{\gamma_2} S_{\gamma_1} (\mathbf{\Lambda} + \mathbf{R}), \gamma_i)/(\gamma_i, \gamma_i) \in \mathbb{N}, \quad i = 1,2,\ldots,k, \end{array} \right. \tag{29}$$

where S_{γ_i} is the Weyl reflection for γ_i, $S_{\gamma_0} \equiv 1$, and $\mathbf{R}$ is the half sum of all positive roots. Eq. (29) gives the weights of all possible extremal vectors of $d_{\mathbf{\Lambda}}$. Gruber *et al.*[44] put forward an algorithm to construct all extremal vectors. Suppose that first one knows an extremal vector $Z_1 \in \Omega_-$, which defines an ideal $\Omega_- Z_1 \subset \Omega_-$. Let $\mathbf{M}_1$ denote the weight of Z_1, then $\mathbf{M}_1$ is the highest weight of a new elementary representation $d_{\mathbf{M}_1}$ defined on $\Omega_- Z_1$. If $d_{\mathbf{M}_1}$ contains another extremal vector $Z_2 \in \Omega_- Z_1$ with the weight $\mathbf{M}_2$, one can construct another elementary representation $d_{\mathbf{M}_2}$ on the ideal $\Omega_- Z_2 Z_1 \subset \Omega_- Z_1$, with $\mathbf{M}_2$ being its highest weight. $\Omega_- Z_2 Z_1$ is also an ideal of Ω_-, and therefore the vector $Z_2 Z_1$ is the extremal vector defining $\Omega_- Z_2 Z_1$ with respect to Ω_-. Repeating this procedure, one may find the ith extremal vector $Y_i \equiv Z_i Z_{i-1} \ldots Z_1$, and the corresponding ideal $I_{Y_i} \equiv \Omega_- Z_i Z_{i-1} \ldots Z_1$. Thus, all ideals form an ideal chain $I_{Y_i} \subset I_{Y_{i-1}} \subset \cdots \subset I_{Y_1} \subset \Omega_-$ naturally.

Now consider an elementary representation $d_{\mathbf{\Lambda}}$ of G_2, with $\mathbf{\Lambda}$ being dominant, i.e., $\langle \mathbf{\Lambda}, \alpha_i \rangle \equiv 2(\mathbf{\Lambda}, \alpha_i)/(\alpha_i, \alpha_i)$ are nonnegative integers for both simple roots α_1 and α_6. From the explicit matrix elements of $d_{\mathbf{\Lambda}}$ given by Eq. (26), one can find that for the simple root vectors $E_{-1}^{m_1}$ and $E_{-6}^{m_6}$, most terms in Eq. (26) vanish, and the second condition of Eq. (28) becomes

$$\left\{ \begin{array}{ll} \rho(E_1)E_{-1}^{m_1} & = m_1 \left(\frac{\Lambda_1}{4} - \frac{\sqrt{3}\Lambda_2}{4} - \frac{m_1 - 1}{8} \right) E_{-1}^{m_1 - 1} \\ & = m_1 \frac{(\alpha_1,\alpha_1)}{2} \left[\frac{2(\mathbf{\Lambda},\alpha_1)}{(\alpha_1,\alpha_1)} - (m_1 - 1) \right] E_{-1}^{m_1 - 1} = 0, \\ \rho(E_i)E_{-1}^{m_1} & = 0, \quad i = 2,3,\ldots,6 \end{array} \right. \tag{30}$$

for $E_{-1}^{m_1}$, and

$$\left\{\begin{array}{ll} \rho(E_i)E_{-6}^{m_6} & = 0, \quad i = 1, 2, \ldots, 5, \\ \rho(E_6)E_{-6}^{m_6} & = m_6(\frac{\Lambda_2}{2\sqrt{3}} - \frac{m_6-1}{24})E_{-6}^{m_6-1} \\ & = m_6\frac{(\alpha_6,\alpha_6)}{2}\left[\frac{2(\mathbf{\Lambda},\alpha_6)}{(\alpha_6,\alpha_6)} - (m_6-1)\right]E_{-6}^{m_6-1} = 0 \end{array}\right. \tag{31}$$

for $E_{-6}^{m_6}$. Solving (30) and (31) gives

$$\left\{\begin{array}{l} m_1 = \frac{2(\mathbf{\Lambda},\alpha_1)}{(\alpha_1,\alpha_1)} + 1 = p + 1 = \frac{2(\mathbf{\Lambda}+\mathbf{R},\alpha_1)}{(\alpha_1,\alpha_1)} \equiv P, \\ m_6 = \frac{2(\mathbf{\Lambda},\alpha_6)}{(\alpha_6,\alpha_6)} + 1 = q + 1 = \frac{2(\mathbf{\Lambda}+\mathbf{R},\alpha_6)}{(\alpha_6,\alpha_6)} \equiv Q. \end{array}\right. \tag{32}$$

Since $\mathbf{\Lambda}$ is dominant, p, q are nonnegative integers and P, Q are positive integers.

Let Y_{01} denote the extremal vector $\mathbf{1}$. Here we use the double-subscript because there are two series of extremal vectors, and therefore two ideal chains (as we can see below). The first subscript is used to label the layers, and the second one the ideal chains. It follows from Eqs. (30) - (32) that we obtain in Ω_- two extremal vectors $Y_{11} = E_{-1}^{P}$ and $Y_{12} = E_{-6}^{Q}$, with their respective weights being

$$\mathbf{M}_{11} = \mathbf{\Lambda} - P\alpha_1 = \mathbf{\Lambda} - \frac{2(\mathbf{\Lambda}+\mathbf{R},\alpha_1)}{(\alpha_1,\alpha_1)}\alpha_1 = S_{\alpha_1}(\mathbf{\Lambda}+\mathbf{R}) - \mathbf{R}, \tag{33}$$

$$\mathbf{M}_{12} = \mathbf{\Lambda} - Q\alpha_6 = \mathbf{\Lambda} - \frac{2(\mathbf{\Lambda}+\mathbf{R},\alpha_6)}{(\alpha_6,\alpha_6)}\alpha_6 = S_{\alpha_6}(\mathbf{\Lambda}+\mathbf{R}) - \mathbf{R}. \tag{34}$$

Y_{11} and Y_{12} are the first layer of extremal vectors below $\mathbf{1}$ (see Fig. 3).

Now we use Y_{11} and $\mathbf{M}_{11}$ in substitution for Y_{01} and $\mathbf{\Lambda}$ respectively, then Eqs. (30) - (34) give the extremal vector $Z_{21} = E_{-6}^{Q'}$ in $\Omega_- Y_{11}$, and Eq. (32) gives $P' = 3P+Q$. The corresponding extremal vector with respect to Ω_- is therefore $Y_{21} = E_{-6}^{Q'}Y_{11} = E_{-6}^{3P+Q}E_{-1}^{P}$. Similar arguments for Y_{12} give another extremal vector $Y_{22} = E_{-1}^{P'}Y_{12} = E_{-1}^{P+Q}E_{-6}^{Q}$. Obviously, Y_{21} and Y_{22} are products of two simple root vectors, and they are the second layer of extremal vectors below $\mathbf{1}$.

Repeating this procedure, we can obtain all twelve extremal vectors of G_2 as follows:

$$\begin{array}{cc} \multicolumn{2}{c}{Y_{01} \equiv \mathbf{1},} \\ Y_{11} = E_{-1}^{P}Y_{01}, & Y_{12} = E_{-6}^{Q}Y_{01}, \\ Y_{21} = E_{-6}^{3P+Q}Y_{11}, & Y_{22} = E_{-1}^{P+Q}Y_{12}, \\ Y_{31} = E_{-1}^{2P+Q}Y_{21}, & Y_{32} = E_{-6}^{3P+2Q}Y_{22}, \\ Y_{41} = E_{-6}^{3P+2Q}Y_{31}, & Y_{42} = E_{-1}^{2P+Q}Y_{32}, \\ Y_{51} = E_{-1}^{P+Q}Y_{41}, & Y_{52} = E_{-6}^{3P+Q}Y_{42}, \\ \multicolumn{2}{c}{Y_{61} = E_{-6}^{Q}Y_{51} = E_{-1}^{P}Y_{52}.} \end{array} \tag{35}$$

Although here we start from the highest extremal vector $\mathbf{1}$, it is easy to verify that starting from any extremal vector, we may recover the same twelve extremal vectors. Fig. 3 shows the positions of these twelve extremal vectors on the weight diagram of $d_{\mathbf{\Lambda}}$. Y_{ik} $(i = 1, 2, \ldots, 6, k = 1, 2)$ belong to the ith layer below $\mathbf{1}$, which are products of i simple root vectors. Especially, for the 6th layer there is only one extremal vector Y_{61}, because its two forms in Eq. (35) are in fact equal (This kind of equivalence has been generally proved by Verma[46]). Each of these extremal vectors defines an ideal $I_{ik} \equiv \Omega_- Y_{ik}$ of Ω_-, which carries an elementary representation $d_{\mathbf{M}_{ik}}$.

If some extremal vector Y_{ik} belongs to the higher layer than Y_{jl} (i.e. $i < j$), then its weight $\mathbf{M}_{ik}$ is higher than the weight $\mathbf{M}_{jl}$ of Y_{jl}, and their corresponding ideals satisfy $I_{jl} \subset I_{ik}$, which can be shown by BGG theorem, therefore, we have $I_{ik} + I_{jl} = I_{ik}$. In order to find new ideals with respect to an elementary representation, we need turn to the sums of subspaces defined by the extremal vectors belonging to the same layer

$$I_{j1} + I_{j2}, \quad j = 1, 2, \ldots, 5. \tag{36}$$

These five subspaces, which have no highest weight vectors, are new ideals of spaces I_{ik} $(i < j)$.

Using these extremal vectors and ideals, one can investigate various new representations on the quotient spaces of I_{ik}, i.e., on I_{ik}/I_{jl} $(i < j)$ and $I_{ik}/(I_{j1} + I_{j2})$ $(i < j)$. If their matrix elements can be explicitly determined, then one may obtain new oscillator realizations of G_2.

V.2 Representations on I_{ik}/I_{jl} and a new five-boson realization

Since all spaces I_{jl} with $j > i$ are ideals of I_{ik}, it is easy to determine that there are 61 different quotient spaces I_{ik}/I_{jl} in all. We denote by $d_{I_{ik}/I_{jl}}$ the representation on



Figure 3: The twelve extremal vectors Y_{ik} on the weight diagram of the elementary representation $d_{\mathbf{\Lambda}}$. $Y_{01} \equiv \mathbf{1}$ is the highest weight vector of $d_{\mathbf{\Lambda}}$. Extremal vectors below $\mathbf{1}$ can be categorized into different layers: Y_{ik} $(i = 1, 2, \ldots, 6)$ belong to the ith layer below $\mathbf{1}$. Extremal vectors in the same layer are not higher nor lower than each other, while Y_{ik} is always higher than Y_{jl} if $i < j$.

the quotient space I_{ik}/I_{jl}. Except that the extremal vectors Y_{51} and Y_{52} have only one extremal vector (Y_{61}) below them, the other extremal vectors Y_{ik} $(i \leq 4, k = 1, 2)$ have more than two different extremal vectors below them. Therefore, $d_{I_{51}/I_{61}}$ and $d_{I_{52}/I_{61}}$ are infinite-dimensional irreducible, while the other 59 representations $d_{I_{ik}/I_{jl}}$ $(i \leq 4, i < j)$ are infinite-dimensional indecomposable.

Matrix elements of the representation $d_{I_{ik}/I_{jl}}$ $(i < j)$ can be obtained by imposing $\Omega_{-}Y_{jl} = 0$ in $d_{\mathbf{M}_{ik}}$. Note that the extremal vectors given by Eq. (35) are not in the standard order (see Eq. (16)). To get the basis of I_{ik}/I_{jl} explicitly, one must reexpress the equations $\Omega_{-}Y_{jl} = 0$ into the standard ordered form by making use of Eqs. (17)-(19). For the general case, this re-ordering is too complicated to be done and unenlightening. However, for the special case $d_{I_{01}/I_{12}}$, the equations $\Omega_{-}Y_{12} = 0$ are already in the standard order, therefore the explicit matrix elements can be simply obtained from $d_{\mathbf{\Lambda}}$ by setting $E_{-1}^{m_1}E_{-2}^{m_2}E_{-3}^{m_3}E_{-4}^{m_4}E_{-5}^{m_5}E_{-6}^{Q+m_6} = 0$ for all $m_i \in \mathbb{Z}^+$. Thus, the representation space of $d_{I_{01}/I_{12}}$ is

$$V_{I_{01}/I_{12}} : \{E_{-1}^{m_1}E_{-2}^{m_2}E_{-3}^{m_3}E_{-4}^{m_4}E_{-5}^{m_5}E_{-6}^{m_6} | m_6 < Q, m_i \in \mathbb{Z}^+\}. \tag{37}$$

If $q = 2(\Lambda, \alpha_6)/(\alpha_6, \alpha_6) = 0$, i.e., $\Lambda_2 = 0$, then $Q = 1$, $V_{I_{01}/I_{12}}$ becomes

$$V'_{I_{01}/I_{12}} : \{E_{-1}^{m_1}E_{-2}^{m_2}E_{-3}^{m_3}E_{-4}^{m_4}E_{-5}^{m_5} | m_i \in \mathbb{Z}^+\}. \tag{38}$$

By discarding all terms including m_6 in $d_{\mathbf{\Lambda}}$ given by Eq. (26), one may get the explicit matrix elements of the representation $d_{I_{01}/I_{12}}$, which are not given here. Furthermore, with the help of Eq. (10), one may obtain an inhomogeneous five-boson realization of G_2:

$$\begin{aligned}
B(E_{-1}) &= a_1^\dagger, \\
B(E_{-2}) &= a_2^\dagger, \\
B(E_{-3}) &= a_3^\dagger, \\
B(E_{-4}) &= a_4^\dagger - \frac{1}{2\sqrt{2}}a_3^\dagger a_2, \\
B(E_{-5}) &= a_5^\dagger + \frac{1}{2\sqrt{2}}a_3^\dagger a_1, \\
B(E_{-6}) &= -\frac{1}{2\sqrt{2}}a_2^\dagger a_1 - \frac{1}{\sqrt{6}}a_4^\dagger a_2 - \frac{1}{8\sqrt{3}}a_3^\dagger a_2^2 - \frac{1}{2\sqrt{2}}a_5^\dagger a_4, \\
B(E_1) &= \frac{1}{48\sqrt{6}}a_3^\dagger a_2^3 - \frac{1}{8\sqrt{3}}a_4^\dagger a_2^2 - \frac{1}{8}a_5^\dagger a_2 a_4 - \frac{1}{2\sqrt{2}}a_5^\dagger a_3 \\
&\quad + \frac{1}{4}\left[\Lambda_1 - \frac{1}{2}\left(\hat{n}_1 + \hat{n}_2 + \hat{n}_3 - \hat{n}_5\right)\right] a_1, \\
B(E_2) &= \frac{1}{2\sqrt{2}}a_4^\dagger a_3 + \frac{1}{16\sqrt{6}}a_3^\dagger a_1 a_4^2 - \frac{1}{4\sqrt{3}}a_2^\dagger a_1 a_4 - \frac{1}{8}a_4^\dagger a_1 a_5 \\
&\quad + \frac{1}{4}\left[\Lambda_1 - \frac{1}{6}\left(3\hat{n}_1 + 2\hat{n}_2 + 3\hat{n}_3 + \hat{n}_4\right)\right] a_2 - \frac{1}{8\sqrt{3}}a_5^\dagger a_4^2, \\
B(E_3) &= -\frac{1}{48\sqrt{6}}a_1^\dagger a_2^3 + \frac{1}{8\sqrt{2}}\left[\Lambda_1 - \frac{1}{6}\left(2\hat{n}_2 + \hat{n}_4 + 3\hat{n}_5\right)\right] a_2 a_4 - \frac{1}{16\sqrt{6}}a_4^\dagger a_2^2 a_5
\end{aligned}$$

$$
\begin{aligned}
& +\frac{1}{192}a_3^\dagger a_2^2 a_4^2 + \frac{1}{2}\left[\Lambda_1 - \frac{1}{4}\left(\hat{n}_1+\hat{n}_2+\hat{n}_3+\hat{n}_4+\hat{n}_5\right)\right] a_3 \\
& +\frac{1}{48\sqrt{6}}a_5^\dagger a_4^3 + \frac{1}{16\sqrt{6}}a_2^\dagger a_1 a_4^2 - \frac{1}{96\sqrt{3}}a_3^\dagger a_1 a_4^3 \\
& -\frac{1}{8\sqrt{2}}\left[\Lambda_1 - \frac{1}{2}(\hat{n}_4+\hat{n}_5)\right] a_1 a_5, \\
B(E_4) \;=\; & -\frac{1}{8\sqrt{3}}a_1^\dagger a_2^2 + \frac{1}{24\sqrt{2}}a_3^\dagger a_2 a_4^2 - \frac{1}{4\sqrt{3}}a_4^\dagger a_2 a_5 \\
& -\frac{1}{2\sqrt{2}}a_2^\dagger a_3 + \frac{1}{4}\left[\Lambda_1 - \frac{1}{6}\left(4\hat{n}_2+\hat{n}_4+3\hat{n}_5\right)\right] a_4, \\
B(E_5) \;=\; & \frac{1}{2\sqrt{2}}a_1^\dagger a_3 - \frac{1}{8\sqrt{3}}a_2^\dagger a_4^2 + \frac{1}{24\sqrt{6}}a_3^\dagger a_4^3 + \frac{1}{4}\left[\Lambda_1 - \frac{1}{2}(\hat{n}_4+\hat{n}_5)\right] a_5, \\
B(E_6) \;=\; & -\frac{1}{2\sqrt{2}}a_1^\dagger a_2 - \frac{1}{\sqrt{6}}a_2^\dagger a_4 - \frac{1}{2\sqrt{2}}a_4^\dagger a_5 + \frac{1}{8\sqrt{3}}a_3^\dagger a_4^2, \\
B(H_1) \;=\; & \Lambda_1 - \frac{1}{4}\left(\hat{n}_1+\hat{n}_2+2\hat{n}_3+\hat{n}_4+\hat{n}_5\right), \\
B(H_2) \;=\; & \frac{1}{4\sqrt{3}}\left(3\hat{n}_1+\hat{n}_2-\hat{n}_4-3\hat{n}_5\right).
\end{aligned}
$$

If one chooses another basis of Ω_- in a different order with E_{-1} on the right (for example, $E_{-6}^{m_6}E_{-5}^{m_5}\ldots E_{-1}^{m_1}$) instead of the standard order in Eq. (24), then using the same method one can get another indecomposable representation $d_{I_{01}/I_{11}}$ involving five generators E_{-i} $(i=2,\ldots,6)$, and correspondingly, one can obtain a new inhomogeneous five-boson realization.

V.3 Representations on the quotient spaces $I_{ik}/(I_{j1}+I_{j2})$ and a three-fermion realization of fundamental representation $(0,1)$

We denote by $d_{I_{ik}/(I_{j1}+I_{j2})}$ the representation on the quotient space $I_{ik}/(I_{j1}+I_{j2})$. Owing to the fact that the spaces $I_{j1}+I_{j2}$ $(j=1,2,\ldots,5)$ are ideals of the spaces I_{ik} with $i<j$, there are in all 25 different quotient spaces $I_{ik}/(I_{j1}+I_{j2})$. For the nine cases of the nearest-neighbor layers the representations $d_{I_{j-1,k}/(I_{j1}+I_{j2})}$ are irreducible, because $I_{j1}+I_{j2}$ are the largest ideal of $I_{j-1,k}$. For the remaining cases the corresponding representations $d_{I_{ik}/(I_{j1}+I_{j2})}$ $(i<j-1)$ are indecomposable. These representations are infinite-dimensional except for $d_{I_{01}/(I_{11}+I_{12})}$.

The matrix elements of $d_{I_{ik}/(I_{j1}+I_{j2})}$ may be determined from $d_{\mathbf{M}_{ik}}$ by setting in I_{ik}:

$$
\left\{\begin{array}{l}
\Omega_- Y_{j1}=0, \\
\Omega_- Y_{j2}=0.
\end{array}\right. \tag{39}
$$

For the infinite-dimensional representations $d_{I_{ik}/(I_{j1}+I_{j2})}$ $(i\neq 0)$, the PBW basis of the subspaces $\Omega_- Y_{jl}$ $(l=1,2)$ here are not in the standard order, however, the re-ordering is too complicated to be done. Whereas, for $d_{I_{01}/(I_{11}+I_{12})}$ the re-ordering becomes possible since it is finite-dimensional. The representation $d_{I_{01}/(I_{11}+I_{12})}$ can also be labeled by Dynkin symbol (p,q) (where $p = P-1$, $q = Q-1$), which is related to $\mathbf{\Lambda}$ by Eq. (32). As an example, we now calculate the simplest case of the fundamental representation $(0,1)$, which is seven-dimensional.



Figure 4: Dots are the seven weights (μ_i, $i = 1,\ldots,7$) of the weight diagram of the fundamental representation (0,1) of G_2, and dim $V(\mu_i) = 1$. Circles denote the weights of the other weight spaces of $d_{\mathbf{\Lambda}}$ (here $\mathbf{\Lambda} = (\frac{1}{4}, \frac{1}{4\sqrt{3}})$), which are void in the quotient space $I_{01}/(I_{11}+I_{12})$. I_{01} is the whole representation space of $d_{\mathbf{\Lambda}}$, $I_{11} = \Omega_- E_{-1}$ and $I_{12} = \Omega_- E_{-6}^2$.

For (0,1), it follows from Eq. (32) that $\mathbf{\Lambda}$ is $(\frac{1}{4}, \frac{1}{4\sqrt{3}})$ and $P = 1, Q = 2$, i.e., $Y_{11} = E_{-1}^{1}, Y_{12} = E_{-6}^{2}$. The weight diagram of $(0,1)$ is shown in Fig. 4. We denote by μ_i $(i = 1, 2, \ldots, 7)$ the seven weights of $(0,1)$, and by $V(\mu_i)$ and $\Omega_-(\mu_i)$ the corresponding weight spaces of $(0,1)$ and $d_{Y_{01}}$ respectively. The standard ordered PBW bases of $\Omega_-(\mu_i)$ read respectively

$$\begin{aligned}
\Omega_-(\mu_1):&\quad \{\mathbf{1}\},\\
\Omega_-(\mu_2):&\quad \{E_{-6}\},\\
\Omega_-(\mu_3):&\quad \{E_{-2}, E_{-1}E_{-6}\},\\
\Omega_-(\mu_4):&\quad \{E_{-4}, E_{-2}E_{-6}, E_{-1}E_{-6}^2\},\\
\Omega_-(\mu_5):&\quad \{E_{-5}, E_{-4}E_{-6}, E_{-2}E_{-6}^2, E_{-1}E_{-6}^3\},\\
\Omega_-(\mu_6):&\quad \{E_{-3}, E_{-2}E_{-4}, E_{-1}E_{-5}, E_{-2}^2E_{-6}, E_{-1}E_{-4}E_{-6}, E_{-1}E_{-2}E_{-6}^2, E_{-1}^2E_{-6}^3\},\\
\Omega_-(\mu_7):&\quad \big\{E_{-4}^2, E_{-2}E_{-5}, E_{-3}E_{-6}, E_{-2}E_{-4}E_{-6}, E_{-1}E_{-5}E_{-6},\\
&\quad E_{-2}^2E_{-6}^2, E_{-1}E_{-4}E_{-6}^2, E_{-1}E_{-2}E_{-6}^3, E_{-1}^2E_{-6}^3\big\}\,. \qquad (40)
\end{aligned}$$

On weight spaces $\Omega_-(\mu_1)$ and $\Omega_-(\mu_2)$, the equations

$$\left\{\begin{array}{l}
\Omega_- Y_{12} = \{E_{-1}^{m_1}E_{-2}^{m_2}E_{-3}^{m_3}E_{-4}^{m_4}E_{-5}^{m_5}E_{-6}^{m_6}\}E_{-6}^2 = 0,\\
\Omega_- Y_{11} = \{E_{-1}^{m_1}E_{-2}^{m_2}E_{-3}^{m_3}E_{-4}^{m_4}E_{-5}^{m_5}E_{-6}^{m_6}\}E_{-1} = 0
\end{array}\right. \qquad (41)$$

give no constraint relations, so that we get $V(\mu_1) = \Omega_-(\mu_1)$ and $V(\mu_2) = \Omega_-(\mu_2)$. On $\Omega_-(\mu_3)$, the condition $\Omega_- Y_{12} = 0$ gives no constraints, while $\Omega_- Y_{11} = 0$ requires $E_{-6}Y_{11} = E_{-6}E_{-1} = 0$, i.e., $E_{-1}E_{-6} - 1/(2\sqrt{2})E_{-2} = 0$, which shows that the two basis elements $E_{-1}E_{-6}$ and E_{-2} are linearly dependent, therefore, for $(0,1)$

the corresponding weight space is $V(\mu_3) = \{E_{-2} = 2\sqrt{2}E_{-1}E_{-6}\}$, which is one-dimensional. It follows from Eq. (41) that by the same method we can determine the constraint relations among basis elements for $\Omega_-(\mu_4), \Omega_-(\mu_5), \Omega_-(\mu_6)$, and $\Omega_-(\mu_7)$. Thus the bases of the seven weight spaces $V(\mu_i)$ of $(0,1)$ are respectively

$$\begin{array}{ll}
V(\mu_1): & \{\mathbf{1}\}, \\
V(\mu_2): & \{E_{-6}\}, \\
V(\mu_3): & \{E_{-2} = 2\sqrt{2}E_{-1}E_{-6}\}, \\
V(\mu_4): & \{E_{-4} = 2\sqrt{6}E_{-2}E_{-6}\}, \\
V(\mu_5): & \{E_{-4}E_{-6} = \frac{1}{6\sqrt{2}}E_{-5}\}, \\
V(\mu_6): & \{E_{-2}E_{-4} = 3\sqrt{2}E_{-3} = -\frac{2}{3}E_{-1}E_{-5} = 2\sqrt{6}E_{-2}^2E_{-6} = -4\sqrt{2}E_{-1}E_{-4}E_{-6}\}, \\
V(\mu_7): & \{E_{-2}E_{-4}E_{-6} = -\frac{1}{4\sqrt{6}}E_{-4}^2 = \frac{1}{6\sqrt{2}}E_{-2}E_{-5} = \frac{1}{6\sqrt{2}}E_{-3}E_{-6}\}.
\end{array} \tag{42}$$

All of them are one-dimensional.

Eq. (42) shows that for (0,1) if we choose the basis

$$V_{(0,1)} : \{\mathbf{1}, E_{-6}, E_{-2}, E_{-4}, E_{-4}E_{-6}, E_{-2}E_{-4}, E_{-2}E_{-4}E_{-6}\}, \tag{43}$$

then we get the representation space with its PBW basis consisting of only three independent generators with powers no larger than one. This makes it possible to construct a three-fermion realization with respect to (0,1).

The basis elements of $V_{(0,1)}$ given by Eq. (43) may be written in the convenient form

$$\{X_{(m_2,m_4,m_6)} = E_2^{m_2}E_4^{m_4}E_6^{m_6} | m_2, m_4, m_6 = 0, 1\}.$$

The nonzero actions of $E_{\pm i}$ $(i = 1, \ldots, 6)$ upon the basis elements in Eq. (43) can be obtained by Eqs. (13) - (14). Using the method very parallel to the bosonic case (see Eqs. (3) - (7)), the fermion Fock space

$$\mathcal{F}_f : \{|m_2, m_4, m_6\rangle = (f_2^\dagger)^{m_2}(f_4^\dagger)^{m_4}(f_6^\dagger)^{m_6}|0\rangle\}$$

may be defined, where $f_i^\dagger$ and f_i $(i = 2, 4, 6)$ are the creation and annihilation operators of the ith fermion, and $|0\rangle$ denotes the vacuum state of $\mathcal{F}_f$. The anti-commutation relations are

$$\{f_i, f_i^\dagger\} = \delta_{ij}, \quad \{f_i, f_j\} = \{f_i^\dagger, f_j^\dagger\} = 0. \tag{44}$$

The following relations may be used to simplify calculations

$$\left\{ \begin{array}{l}
\hat{n}_i^f = f_i^\dagger f_i = (f_i^\dagger f_i)^2 = (\hat{n}_i^f)^2, \\
(1 - \hat{n}_i^f)^2 = 1 - \hat{n}_i^f, \\
f_i f_i^\dagger f_i = (1 - f_i^\dagger f_i) f_i = f_i, \\
f_i^\dagger f_i f_i^\dagger = f_i^\dagger (1 - f_i^\dagger f_i) = f_i^\dagger,
\end{array} \right. \tag{45}$$

where $\hat{n}_i^f$ is the particle number operator of the ith fermion.

The actions of E_{-1} on the basis of $V_{(0,1)}$ are

$$\left\{\begin{array}{l} \phi(E_{-1})X_{(0,0,1)} = \frac{1}{2\sqrt{2}}X_{(1,0,0)}, \\ \phi(E_{-1})X_{(0,1,1)} = -\frac{1}{4\sqrt{2}}X_{(1,1,0)}, \\ \phi(E_{-1})X_{(m_2,m_4,m_6)} = 0 \quad \text{for other five basis elements.} \end{array}\right. \tag{46}$$

By making use of the δ-function, Eq. (46) can be written in a more compact form

$$\phi(E_{-1})X_{(m_2,m_4,m_6)} = (\frac{1}{2\sqrt{2}}\delta_{0,m_2}\delta_{0,m_4}\delta_{1,m_6} - \frac{1}{4\sqrt{2}}\delta_{0,m_2}\delta_{1,m_4}\delta_{1,m_6})X_{(m_2+1,m_4,m_6-1)}. \tag{47}$$

Since m_i can take 0 or 1 only, all terms involving $\delta_{0,m_i}X_{(m_i-1)}$ and $\delta_{1,m_i}X_{(m_i+1)}$ vanish. For the non-zero terms there exist the following correspondence relations:

$$\left\{\begin{array}{l} \delta_{0,m_i}X_{(m_i+1)} \longmapsto f_i^\dagger(1-f_i^\dagger f_i)|m_i\rangle = f_i^\dagger|m_i\rangle, \\ \delta_{0,m_i}X_{(m_i)} \longmapsto (1-f_i^\dagger f_i)|m_i\rangle = (1-\hat{n}_i^f)|m_i\rangle, \\ \delta_{1,m_i}X_{(m_i)} \longmapsto f_i^\dagger f_i|m_i\rangle = \hat{n}_i^f|m_i\rangle, \\ \delta_{1,m_i}X_{(m_i-1)} \longmapsto f_i(f_i^\dagger f_i)|m_i\rangle = f_i|m_i\rangle. \end{array}\right. \tag{48}$$

Then from Eq. (47) one gets the fermion realization of E_{-1}

$$F(E_{-1}) = \frac{1}{2\sqrt{2}}f_2^\dagger(1-f_4^\dagger f_4)f_6 + \frac{1}{4\sqrt{2}}f_2^\dagger f_4^\dagger f_4 f_6 = \frac{1}{2\sqrt{2}}(1-\frac{1}{2}\hat{n}_4^f)f_2^\dagger f_6. \tag{49}$$

Realizations of the remaining generators of G_2 can be calculated by the same method. They are

$$\begin{array}{rcl} F(E_{-2}) &=& (1+\hat{n}_4^f\hat{n}_6^f-\hat{n}_6^f)f_2^\dagger + \frac{1}{2\sqrt{6}}(1-\hat{n}_2^f)f_4^\dagger f_6, \\ F(E_{-3}) &=& 3\sqrt{2}(1+\hat{n}_6^f)f_2^\dagger f_4^\dagger, \\ F(E_{-4}) &=& (1-\frac{1}{2}\hat{n}_2^f-\frac{1}{2}\hat{n}_2^f\hat{n}_6^f)f_4^\dagger + 4\sqrt{6}\hat{n}_4^f f_2^\dagger f_6^\dagger, \\ F(E_{-5}) &=& 6\sqrt{2}f_4^\dagger f_6^\dagger, \\ F(E_{-6}) &=& (1-\hat{n}_2^f+\hat{n}_4^f+\hat{n}_2^f\hat{n}_4^f)f_6^\dagger + \frac{1}{2\sqrt{6}}(1-\hat{n}_6^f)f_2 f_4^\dagger, \\ F(E_1) &=& -\frac{1}{2\sqrt{2}}(1+\hat{n}_4^f)f_2 f_6^\dagger, \\ F(E_2) &=& \frac{1}{24}(1+\hat{n}_4^f-\hat{n}_6^f)f_2 - \frac{1}{\sqrt{6}}(1-\hat{n}_2^f)f_4 f_6^\dagger, \\ F(E_3) &=& -\frac{1}{24\sqrt{2}}(1-\frac{1}{2}\hat{n}_6^f)f_2 f_4, \\ F(E_4) &=& \frac{1}{12}(1-\frac{1}{2}\hat{n}_6^f-\frac{1}{2}\hat{n}_2^f\hat{n}_6^f)f_4 - \frac{1}{48\sqrt{6}}\hat{n}_4^f f_2 f_6, \\ F(E_5) &=& -\frac{1}{48\sqrt{2}}f_4 f_6, \end{array}$$

$$
\begin{aligned}
F(E_6) &= \frac{1}{24}(1-\hat{n}_2^f+\frac{1}{2}\hat{n}_2^f\hat{n}_4^f)f_6-\frac{1}{\sqrt{6}}(1-\hat{n}_6^f)f_2^\dagger f_4, \\
F(H_1) &= \frac{1}{4}(1-\hat{n}_2^f-\hat{n}_4^f), \\
F(H_2) &= \frac{1}{4\sqrt{3}}(1+\hat{n}_2^f-\hat{n}_4^f-2\hat{n}_6^f). \qquad (50)
\end{aligned}
$$

It is easy to verify with the help of Eqs. (44) and (45) that they satisfy the commutation relations of G_2. Hence from the fundamental representation $(0,1)$, we obtain a new three-fermion realization, which is different from the fermion realizations based upon two Lie algebra chains $so(8)\supset G_2$ and $G_2\supset su(3)$.[47]

VI SUMMARY AND DISCUSSIONS

In this paper we have obtained the explicit matrix elements of the master representation ρ of G_2, which is defined on the space Ω of the universal enveloping algebra of G_2, and discussed various representations induced from ρ, such as the representations on the invariant subspaces of Ω and the representations on the quotient spaces with respect to these invariant subspaces. Particularly, we have investigated in detail the elementary representations of G_2, which are defined on the subspace Ω_- (generated by E_{-i} only) of Ω and have an highest weight, and from which we have realized G_2 by six boson pairs. For the elementary representation with a fixed dominate $\mathbf{\Lambda}$ being its highest weight, we have obtained all twelve extremal vectors in explicit form, each of them defines an ideal I_{ik} of Ω_-, which carries an elementary sub-representation. With the help of these extremal vectors we have investigated the representations on the quotient spaces of Ω_- with fewer generators, and consequently constructed the five-boson realization from $d_{I_{01}/I_{12}}$ and the three-fermion realization from the special case of $d_{I_{01}/(I_{11}+I_{12})}=(0,1)$.

If one makes use of the following corresponding relations between the boson operators and the differential operators

$$
a_i^\dagger \Leftrightarrow x_i, \quad a_i \Leftrightarrow \frac{\partial}{\partial x_i},
$$

where x_i and $\frac{\partial}{\partial x_i}$ satisfy the commutation relations

$$
\left[\frac{\partial}{\partial x_i}, x_j\right]=\delta_{ij}, \quad \left[\frac{\partial}{\partial x_i}, \frac{\partial}{\partial x_j}\right]=[x_i,x_j]=0,
$$

then one can obtain immediately the corresponding inhomogeneous differential realizations of G_2 from the six-boson or five-boson realizations obtained in this paper.

If one chooses as the PBW basis for Ω the set of elements in a different order, for example

$$
\Omega:\{E_1^{n_1}E_2^{n_2}E_3^{n_3}E_4^{n_4}E_5^{n_5}E_6^{n_6}E_{-1}^{m_1}E_{-2}^{m_2}E_{-3}^{m_3}E_{-4}^{m_4}E_{-5}^{m_5}E_{-6}^{m_6}H_1^{k_1}H_2^{k_2}\},
$$

then one can study in the same manner the elementary representations on Ω_+ : $\{E_1^{n_1}E_2^{n_2}E_3^{n_3}E_4^{n_4}E_5^{n_5}E_6^{n_6}\}$, the extremal vectors, and the representations on various ideals or quotient spaces with respect to Ω_+. These representations have their respective lowest weights, whereas the representations on the spaces with respect to Ω_- discussed in this paper have the highest weights. Similarly, one can obtain the corresponding boson or fermion realizations.

Similar procedure can also be used to investigate the exceptional Lie superalgebra $G(3)$, which is underway.

ACKNOWLEDGEMENTS

This work is supported by the State Key Basic Research Development Programs (Grant No. 2009CB929402).

A The matrix elements of the master representation ρ of G_2

Here we do not give the concrete values of the coefficients $N_{\alpha,\beta}$ of G_2, thus, if we choose the other basis (for example, the irreducible tensor basis), we can easily get the new master representation by simply replacing the symbols of the basis and the coefficients in ρ. Using the convenient abbreviations: $X_{(mnk)} \equiv X(\ldots, m_i, \ldots)$ and $X_{m_i+k} \equiv X(\ldots, m_i+k, \ldots)$ in Ω, the explicit matrix elements of ρ are

$$
\begin{aligned}
\rho(E_{-1})X_{(mnk)} &= X_{m_1+1}, \\
\rho(E_{-2})X_{(mnk)} &= X_{m_2+1}, \\
\rho(E_{-3})X_{(mnk)} &= X_{m_3+1}, \\
\rho(E_{-4})X_{(mnk)} &= X_{m_4+1} + m_2 N_{-4,-2} X_{m_2-1,m_3+1}, \\
\rho(E_{-5})X_{(mnk)} &= X_{m_5+1} + m_1 N_{-5,-1} X_{m_1-1,m_3+1}, \\
\rho(E_{-6})X_{(mnk)} &= X_{m_6+1} + m_1 N_{-6,-1} X_{m_1-1 m_2+1} + m_2 N_{-6,-2} X_{m_2-1 m_4+1} \\
&\quad + \frac{1}{2} m_2(m_2-1) N_{-6,-2} N_{-4,-2} X_{m_2-1,m_3+1} + m_4 N_{-6,-4} X_{m_4-1 m_5+1}, \\
\rho(E_1)X_{(mnk)} &= X_{n_1+1} \\
&\quad + m_1 \left[\frac{1}{2}(\alpha_1, -\alpha_1)(m_1-1) + \sum_{i=2}^{6} (\alpha_1, -\alpha_i) m_i + \sum_{i=1}^{6} (\alpha_1, \alpha_i) n_i \right] X_{m_1-1} \\
&\quad + m_1 \alpha_1^{(1)} X_{m_1-1,k_1+1} + m_1 \alpha_1^{(2)} X_{m_1-1,k_2+1} \\
&\quad + m_2 N_{1,-2} X_{m_2-1,m_6+1} + \frac{1}{2} m_2(m_2-1) N_{1,-2} N_{-6,-2} X_{m_2-2,m_4+1} \\
&\quad + \frac{1}{6} m_2(m_2-1)(m_2-2) N_{1,-2} N_{-6,-2} N_{-4,-2} X_{m_2-2,m_3+1} \\
&\quad + m_2 m_4 N_{1,-2} N_{-6,-4} X_{m_2-1,m_4-1,m_5+1} \\
&\quad + m_3 N_{1,-3} X_{m_3-1,m_5+1}, \\
\rho(E_2)X_{(mnk)} &= X_{n_2+1} \\
&\quad + m_2 \left[m_1 N_{2,-1} N_{6,-2} + \frac{1}{2}(\alpha_2, -\alpha_2)(m_2-1) + \sum_{i=3}^{6} (\alpha_2, -\alpha_i) m_i \right. \\
&\qquad \left. + \sum_{i=1}^{6} (\alpha_2, \alpha_i) n_i \right] X_{m_2-1} \\
&\quad + m_1 N_{2,-1} X_{m_1-1,n_6+1} \\
&\quad + m_1 m_4 N_{2,-1} N_{6,-4} X_{m_1-1,m_2+1,m_4-1} \\
&\quad + \frac{1}{2} m_1 m_4 (m_4-1) N_{2,-1} N_{6,-4} N_{-4,-2} X_{m_1-1,m_3+1,m_4-2} \\
&\quad + m_1 m_5 N_{2,-1} N_{6,-5} X_{m_1-1,m_4+1,m_5-1} \\
&\quad + m_1 m_6 N_{2,-1} \left[\frac{1}{2}(\alpha_6, -\alpha_6)(m_6-1) + \sum_{i=1}^{6} (\alpha_6, \alpha_i) n_i \right] X_{m_1-1,m_6-1}
\end{aligned}
$$

$$
\begin{aligned}
&+m_1 m_6 N_{2,-1}\alpha_6^{(1)} X_{m_1-1,m_6-1,k_1+1}\\
&+m_1 m_6 N_{2,-1}\alpha_6^{(2)} X_{m_1-1,m_6-1,k_2+1}\\
&+m_1 n_1 N_{2,-1} N_{6,1} X_{m_1-1,n_1-1,n_2+1}\\
&+m_1 n_2 N_{2,-1} N_{6,2} X_{m_1-1,n_2-1,n_4+1}\\
&+\frac{1}{2} m_1 n_2 (n_2-1) N_{2,-1} N_{6,2} N_{4,2} X_{m_1-1,n_2-2,n_3+1}\\
&+m_1 n_4 N_{2,-1} N_{6,4} X_{m_1-1,n_4-1,n_5+1}\\
&+m_2 \alpha_2^{(1)} X_{m_2-1,k_1+1} + m_2 \alpha_2^{(2)} X_{m_2-1,k_2+1}\\
&+m_3 N_{2,-3} X_{m_3-1,m_4+1}\\
&+m_4 N_{2,-4} X_{m_4-1,m_6+1}\\
&+\frac{1}{2} m_4 (m_4-1) N_{2,-4} N_{-6,-4} X_{m_4-2,m_5+1}\\
&+m_6 N_{2,-6} X_{m_6-1,n_1+1},
\end{aligned}
$$

$$
\begin{aligned}
\rho(E_3) X_{(mnk)} = {} & X_{n_3+1}\\
&+m_1 N_{3,-1} X_{m_1-1,n_5+1}\\
&+m_1 m_3 N_{3,-1} N_{5,-3} X_{m_3-1}\\
&+m_1 m_4 N_{3,-1} N_{5,-4} X_{m_1-1,m_4-1,n_6+1}\\
&+\frac{1}{2} m_1 m_4 (m_4-1) N_{3,-1} N_{5,-4} N_{6,-4} X_{m_1-1,m_2+1,m_4-2}\\
&+\frac{1}{3} m_1 m_4 (m_4-1)(m_4-2) N_{3,-1} N_{5,-4} N_{6,-4} N_{-4,-2} X_{m_1-1,m_3+1,m_4-3}\\
&+m_1 m_4 m_5 N_{3,-1} N_{5,-4} N_{6,-5} X_{m_1-1,m_5-1}\\
&+m_1 m_4 m_6 N_{3,-1} N_{5,-4} \left[\frac{1}{2}(\alpha_6,-\alpha_6)(m_6-1) + \sum_{i=1}^{6} (\alpha_6,\alpha_i) n_i\right] X_{m_1-1,m_4-1,m_6-1}\\
&+m_1 m_4 m_6 N_{3,-1} N_{5,-4} \alpha_6^{(1)} X_{m_1-1,m_4-1,m_6-1,k_1+1}\\
&+m_1 m_4 m_6 N_{3,-1} N_{5,-4} \alpha_6^{(2)} X_{m_1-1,m_4-1,m_6-1,k_2+1}\\
&+m_1 m_4 n_1 N_{3,-1} N_{5,-4} N_{6,1} X_{m_1-1,m_4-1,n_1-1,n_2+1}\\
&+m_1 m_4 n_2 N_{3,-1} N_{5,-4} N_{6,2} X_{m_1-1,m_4-1,n_2-1,n_4+1}\\
&+\frac{1}{2} m_1 m_4 n_2 (n_2-1) N_{3,-1} N_{5,-4} N_{6,2} N_{4,2} X_{m_1-1,m_4-1,n_2-2,n_3+1}\\
&+m_1 m_4 n_4 N_{3,-1} N_{5,-4} N_{6,4} X_{m_1-1,m_4-1,n_4-1,n_5+1}\\
&+m_1 m_5 N_{3,-1} \left[\frac{1}{2}(\alpha_5,-\alpha_5)(m_5-1) + (\alpha_5,-\alpha_6) m_6 + \sum_{i=1}^{6} (\alpha_5,\alpha_i) n_i\right] X_{m_1-1,m_5-1}\\
&+m_1 m_5 N_{3,-1} \alpha_5^{(1)} X_{m_1-1,m_5-1,k_1+1} + m_1 m_5 N_{3,-1} \alpha_5^{(2)} X_{m_1-1,m_5-1,k_2+1}\\
&+m_1 m_6 N_{3,-1} N_{5,-6} X_{m_1-1,m_6-1,n_4+1}\\
&+\frac{1}{2} m_1 m_6 (m_6-1) N_{3,-1} N_{5,-6} N_{4,-6} X_{m_1-1,m_6-2,n_2+1}\\
&+\frac{1}{6} m_1 m_6 (m_6-1)(m_6-2) N_{3,-1} N_{5,-6} N_{4,-6} N_{2,-6} X_{m_1-1,m_6-3,n_1+1}
\end{aligned}
$$

$$
\begin{aligned}
&+m_1m_6n_2N_{3,-1}N_{5,-6}N_{4,2}X_{m_1-1,m_6-1,n_2-1,n_3+1}\\
&+m_1n_1N_{3,-1}N_{5,1}X_{m_1-1,n_1-1,n_3+1}\\
&+m_2N_{3,-2}X_{m_2-1,n_4+1}\\
&+\frac{1}{2}m_2(m_2-1)N_{3,-2}N_{4,-2}X_{m_2-2,n_6+1}\\
&+\frac{1}{6}m_2(m_2-1)(m_2-2)N_{3,-2}N_{4,-2}N_{6,-2}X_{m_1+1,m_2-3}\\
&+m_2m_3N_{3,-2}N_{4,-3}X_{m_3-1}\\
&+m_2m_4N_{3,-2}\left[\frac{1}{2}(\alpha_4,-\alpha_4)(m_4-1)+\sum_{i=5}^{6}(\alpha_4,-\alpha_i)m_i\right.\\
&\qquad\qquad\qquad\qquad\left.+\sum_{i=1}^{6}(\alpha_4,\alpha_i)n_i\right]X_{m_2-1,m_4-1}\\
&+m_2m_4N_{3,-2}\alpha_4^{(1)}X_{m_2-1,m_4-1,k_1+1}+m_2m_4N_{3,-2}\alpha_4^{(2)}X_{m_2-1,m_4-1,k_2+1}\\
&+m_2m_5N_{3,-2}N_{4,-5}X_{m_2-1,m_5-1,m_6+1}\\
&+m_2m_6N_{3,-2}N_{4,-6}X_{m_2-1,m_6-1,n_2+1}\\
&+\frac{1}{2}m_2m_6(m_6-1)N_{3,-2}N_{4,-6}N_{2,-6}X_{m_2-1,m_6-2,n_1+1}\\
&+m_2n_2N_{3,-2}N_{4,2}X_{m_2-1,n_2-1,n_3+1}\\
&+\frac{1}{2}m_2(m_2-1)m_4N_{3,-2}N_{4,-2}N_{6,-4}X_{m_2-1,m_4-1}\\
&+\frac{1}{4}m_2(m_2-1)m_4(m_4-1)N_{3,-2}N_{4,-2}N_{6,-4}N_{-4,-2}X_{m_2-2,m_3+1,m_4-2}\\
&+\frac{1}{2}m_2(m_2-1)m_5N_{3,-2}N_{4,-2}N_{6,-5}X_{m_2-2,m_4+1,m_5-1}\\
&+\frac{1}{2}m_2(m_2-1)m_6\left[\frac{1}{2}(\alpha_6,-\alpha_6)(m_6-1)+\sum_{i=1}^{6}(\alpha_6,\alpha_i)n_i\right]X_{m_2-2,m_6-1}\\
&+\frac{1}{2}m_2(m_2-1)m_6N_{3,-2}N_{4,-2}\alpha_6^{(1)}X_{m_2-2,m_6-1,k_1+1}\\
&+\frac{1}{2}m_2(m_2-1)m_6N_{3,-2}N_{4,-2}\alpha_6^{(2)}X_{m_2-2,m_6-1,k_2+1}\\
&+\frac{1}{2}m_2(m_2-1)n_1N_{3,-2}N_{4,-2}N_{6,1}X_{m_2-2,n_1-1,n_2+1}\\
&+\frac{1}{2}m_2(m_2-1)n_2N_{3,-2}N_{4,-2}N_{6,2}X_{m_2-2,n_2-1,n_4+1}\\
&+\frac{1}{4}m_2(m_2-1)n_2(n_2-1)N_{3,-2}N_{4,-2}N_{6,2}N_{4,2}X_{m_2-2,n_2-2,n_3+1}\\
&+\frac{1}{2}m_2(m_2-1)n_4N_{3,-2}N_{4,-2}N_{6,4}X_{m_2-2,n_4-1,n_5+1}\\
&+m_3\left[\frac{1}{2}(\alpha_3,-\alpha_3)(m_3-1)+\sum_{i=4}^{6}(\alpha_3,-\alpha_i)m_i+\sum_{i=1}^{6}(\alpha_3,\alpha_i)n_i\right]X_{m_3-1}\\
&+m_3\alpha_3^{(1)}X_{m_3-1,k_1+1}+m_3\alpha_3^{(2)}X_{m_3-1,k_2+1}
\end{aligned}
$$

$$
\begin{aligned}
&+m_4 N_{3,-4} X_{m_4-1,n_2+1} \\
&+\frac{1}{2} m_4(m_4-1) N_{3,-4} N_{2,-4} X_{m_4-2,m_6+1} \\
&+\frac{1}{6} m_4(m_4-1)(m_4-2) N_{3,-4} N_{2,-4} N_{-6,-4} X_{m_4-2,m_5+1} \\
&+m_4 m_6 N_{3,-4} N_{2,-6} X_{m_4-1,m_6-1,n_1+1} \\
&+m_5 N_{3,-5} X_{m_5-1,n_1+1},
\end{aligned}
$$

$$
\begin{aligned}
\rho(E_4) X_{(mnk)} = {} & X_{n_4+1} \\
&+m_2 N_{4,-2} X_{m_2-1,n_6+1} \\
&+\frac{1}{2} m_2(m_2-1) N_{4,-2} N_{6,-2} X_{m_1+1,m_2-2} \\
&+m_2 m_4 N_{4,-2} N_{6,-4} X_{m_4-1} \\
&+\frac{1}{2} m_2 m_4(m_4-1) N_{4,-2} N_{6,-4} N_{-4,-2} X_{m_2-1,m_3+1,m_4-2} \\
&+m_2 m_5 N_{4,-2} N_{6,-5} X_{m_2-1,m_4+1,m_5-1} \\
&+m_2 m_6 N_{4,-2} \left[\frac{1}{2}(\alpha_6,-\alpha_6)(m_6-1) + \sum_{i=1}^{6} (\alpha_6,\alpha_i) n_i \right] X_{m_2-1,m_6-1} \\
&+m_2 m_6 N_{4,-2} \alpha_6^{(1)} X_{m_2-1,m_6-1,k_1+1} + m_2 m_6 N_{4,-2} \alpha_6^{(2)} X_{m_2-1,m_6-1,k_2+1} \\
&+m_2 n_1 N_{4,-2} N_{6,1} X_{m_2-1,n_1-1,n_2+1} \\
&+m_2 n_2 N_{4,-2} N_{6,2} X_{m_2-1,n_2-1,n_4+1} \\
&+\frac{1}{2} m_2 n_2(n_2-1) N_{4,-2} N_{6,2} N_{4,2} X_{m_2-1,n_2-2,n_3+1} \\
&+m_2 n_4 N_{4,-2} N_{6,4} X_{m_2-1,n_4-1,n_5+1} \\
&+m_3 N_{4,-3} X_{m_2+1,m_3-1} \\
&+m_4 \left[\frac{1}{2}(\alpha_4,-\alpha_4)(m_4-1) + \sum_{i=5}^{6} (\alpha_4,-\alpha_i) m_i + \sum_{i=1}^{6} (\alpha_4,\alpha_i) n_i \right] X_{m_4-1} \\
&+m_4 \alpha_4^{(1)} X_{m_4-1,k_1+1} + m_4 \alpha_4^{(2)} X_{m_4-1,k_2+1} \\
&+m_5 N_{4,-5} X_{m_5-1,m_6+1} \\
&+m_6 N_{4,-6} X_{m_6-1,n_2+1} \\
&+\frac{1}{2} m_6(m_6-1) N_{4,-6} N_{2,-6} X_{m_6-2,n_1+1} \\
&+n_2 N_{4,2} X_{n_2-1,n_3+1},
\end{aligned}
$$

$$
\begin{aligned}
\rho(E_5) X_{(mnk)} = {} & X_{n_5+1} \\
&+m_3 N_{5,-3} X_{m_1+1,m_3-1} \\
&+m_4 N_{5,-4} X_{m_4-1,n_6+1} \\
&+\frac{1}{2} m_4(m_4-1) N_{5,-4} N_{6,-4} X_{m_2+1,m_4-2} \\
&+\frac{1}{3} m_4(m_4-1)(m_4-2) N_{5,-4} N_{6,-4} N_{-4,-2} X_{m_3+1,m_4-3} \\
&+m_4 m_5 N_{5,-4} N_{6,-5} X_{m_5-1}
\end{aligned}
$$

$$
\begin{aligned}
&\quad +m_4 m_6 N_{5,-4}\left[\frac{1}{2}\sum_{i=1}^{6}(\alpha_6,-\alpha_6)(m_6-1)+(\alpha_6,\alpha_i)n_i\right]X_{m_4-1,m_6-1}\\
&\quad +m_4 m_6 N_{5,-4}\alpha_6^{(1)}X_{m_4-1,m_6-1,k_1+1}+m_4 m_6 N_{5,-4}\alpha_6^{(2)}X_{m_4-1,m_6-1,k_2+1}\\
&\quad +m_4 n_1 N_{5,-4}N_{6,1}X_{m_4-1,n_1-1,n_2+1}\\
&\quad +m_4 n_2 N_{5,-4}N_{6,2}X_{m_4-1,n_2-1,n_4+1}\\
&\quad +\frac{1}{2}m_4 n_2(n_2-1)N_{5,-4}N_{6,2}N_{4,2}X_{m_4-1,n_2-2,n_3+1}\\
&\quad +m_4 n_4 N_{5,-4}N_{6,4}X_{m_4-1,n_4-1,n_5+1}\\
&\quad +m_5\left[\frac{1}{2}(\alpha_5,-\alpha_5)(m_5-1)+(\alpha_5,-\alpha_6)m_6+\sum_{i=1}^{6}(\alpha_5,\alpha_i)n_i\right]X_{m_5-1}\\
&\quad +m_5\alpha_5^{(1)}X_{m_5-1,k_1+1}+m_5\alpha_5^{(2)}X_{m_5-1,k_2+1}\\
&\quad +m_6 N_{5,-6}X_{m_6-1,n_4+1}\\
&\quad +\frac{1}{2}m_6(m_6-1)N_{5,-6}N_{4,-6}X_{m_6-2,n_2+1}\\
&\quad +\frac{1}{6}m_6(m_6-1)(m_6-2)N_{5,-6}N_{4,-6}N_{2,-6}X_{m_6-3,n_1+1}\\
&\quad +m_6 n_2 N_{5,-6}N_{4,2}X_{m_6-1,n_2-1,n_3+1}\\
&\quad +n_1 N_{5,1}X_{n_1-1,n_3+1},\\
&\rho(E_6)X_{(mnk)}=X_{n_6+1}\\
&\quad +m_2 N_{6,-2}X_{m_1+1,m_2-1}\\
&\quad +m_4 N_{6,-4}X_{m_2+1,m_4-1}\\
&\quad +\frac{1}{2}m_4(m_4-1)N_{6,-4}N_{-4,-2}X_{m_3+1,m_4-2}\\
&\quad +m_5 N_{6,-5}X_{m_4+1,m_5-1}\\
&\quad +m_6\left[\frac{1}{2}(\alpha_6,-\alpha_6)(m_6-1)+\sum_{i=1}^{6}(\alpha_6,\alpha_i)n_i\right]X_{m_6-1}\\
&\quad +m_6\alpha_6^{(1)}X_{m_6-1,k_1+1}+m_6\alpha_6^{(2)}X_{m_6-1,k_2+1}\\
&\quad +n_1 N_{6,1}X_{n_1-1,n_2+1}\\
&\quad +n_2 N_{6,2}X_{n_2-1,n_4+1}\\
&\quad +\frac{1}{2}n_2(n_2-1)N_{6,2}N_{4,2}X_{n_2-2,n_3+1}\\
&\quad +n_4 N_{6,4}X_{n_4-1,n_5+1},\\
&\rho(H_1)X_{(mnk)}=X_{k_1+1}+\sum_{i=1}^{6}\alpha_i^{(1)}(n_i-m_i)X_{(mnk)},\\
&\rho(H_2)X_{(mnk)}=X_{k_1+1}+\sum_{i=1}^{6}\alpha_i^{(2)}(n_i-m_i)X_{(mnk)}.
\end{aligned}
$$

References

[1] G. Racah, Phys. Rev. **76**, 1352 (1949).

[2] H. A. Jahn, Proc. Roy. Soc. (London) A **201**, 516 (1950).

[3] B. H. Flowers, Proc. Roy. Soc. (London) A **210**, 497 (1952).

[4] D. Ruan and H. Z. Sun, Commun. Theor. Phys. **33**, 221 (2000).

[5] I. Morrison, P. W. Pieruschka, and B. G. Wybourne, J. Math. Phys. **32**, 356 (1991).

[6] G. Dall'Agata and N. Prezas, J. High Energy Phys. **10**, 103 (2005).

[7] B. de Carlos, A. Lukas, and S. Morris, J. High Energy Phys. **12**, 018 (2004).

[8] G. Ferretti, P. Salomonson, and D. Tsimpis, J. High Energy Phys. **3**, 004 (2002).

[9] A. Maas and S. Olejnik, J. High Energy Phys. **2**, 070 (2008).

[10] C. Csaki, C. Grojean, and H. Murayama, Phys. Rev. D **67**, 085012 (2003).

[11] J. Patera, J. Math. Phys. **1**, 3027 (1970).

[12] R. Le Blanc and D. J. Rowe, J. Math. Phys. **29**, 758 (1988).

[13] R. Le Blanc and D. J. Rowe, J. Math. Phys. **29**, 767 (1988).

[14] A. M. Bincer and K. Riesselmann, J. Math. Phys. **34**, 5935 (1993).

[15] R. G. Donnelly, S. J. Lewis, and R. Pervine, Discret. Math. **306**, 1285 (2006).

[16] A. Borowiec, J. Lukierski, V. Lyakhovsky, M. Mozrzymas, and V. N. Tolstoy, J. Math. Phys. **46**, 103502 (2005).

[17] R. Coquereaux, R. Rais, and E. H. Tahri, J. Math. Phys. **51**, 092302 (2010).

[18] U. H. Niederer and L. ORaifeartaigh, Fortschr. Phys. **22**, 111 (1974).

[19] E. Chacón, D. Levi, and M. Moshinsky, J. Math. Phys. **17**, 1919 (1976).

[20] M. Flato and C. Fronsdal, J. Math. Phys. **22**, 1100(1982).

[21] M. Flato and C. Fronsdal, Phys. Lett. B **97**, 236 (1980).

[22] Q. Z. Han, H. Z. Sun, M. Zhang, and D. H. Feng, J. Math. Phys. **26**, 1822 (1985).

[23] M. Lesimple, J. Math. Phys. **39**, 6384 (1998).

[24] K. B. Alkalaev, M. Grigoriev, and I. Yu. Tipunin, arXiv: 0811.3999.

[25] D. P. Zhelobenko, Dokl. Akad. Nauk SSSR, **126**, 635 (1959).

[26] I. M. Gel'fand and V. A. Ponomarev, Russ. Math. Surveys **23**, 1 (1968); translation from Usp. Mat. Nauk **23**, 3 (1968).

[27] B. Gruber and W. C. Henneberger, Nuovo Cimento B **77** 203 (1983).

[28] A. Douglas, J. Math. Phys. **26**, 1822 (2006).

[29] R. Lenczewski and B. Gruber, J. Phys. A **19**, 1 (1986).

[30] H. C. Fu, J. Math. Phys. **32**, 767 (1991).

[31] V. Chari and A. Pressley, *A Guide to Quantum Groups* (Cambridge University Press, Cambridge, 1994).

[32] J. Xiao, Canad. J. Math. **49**, 771 (1997).

[33] P. Casati, S. Minniti, and V. Salari, J. Math. Phys. **51**, 033515 (2010).

[34] B. Gruber and A. U. Klimyk, J. Math. Phys. **19**, 2009 (1978).

[35] B. Gruber and A. U. Klimyk, J. Math. Phys. **25**, 755 (1984).

[36] B. Gruber, A. U. Klimyk, and Y. F. Smirnov, Nuovo Cimento **69**, 97 (1982).

[37] B. Gruber and R. Lenczewski, J. Phys. A **16**, 3703 (1983).

[38] D. Ruan, L. H. Chen, and W. Ruan, J. Math. Phys. **41**, 7839 (2000).

[39] D. Ruan, Y. F. Jia, and W. Ruan, J. Math. Phys. **42**, 2718 (2001).

[40] H. D. Doebner, B. Gruber, and M. Lorente, J. Math. Phys. **30**, 594 (1989).

[41] H. C. Fu and C. P. Sun, J. Math. Phys. **31**, 287 (1990).

[42] D. Ruan, C. Wu, and H. Z. Sun, Commun. Theor. Phys. **40**, 73 (2003).

[43] J. E. Humphreys, *Introductions to Lie Algebras and Representation Theory* (Springer-Verlag, New York, 1972).

[44] M. Lorente and B. Gruber, J. Math. Phys. **29**, 1674 (1984).

[45] I. N. Bemshtein, I. M. Gel'fand, and S. I. Gel'fand, Funkcional. Anal. iPrilozen. **5**, 1 (1971) (in Russian).

[46] D. N. Verma, Bull. Am. Math. Soc. **74**, 160 (1968).

[47] L. Frappat, A. Sciarrino, and P. Sorba, *Dictionary on Lie Algebras and Superalgebras* (Academic Press, San Diego, 2000).